

# Multiple Conclusion Linear Logic: Cut Elimination and more

Harley Eades III<sup>1</sup> and Valeria de Paiva<sup>2</sup>

<sup>1</sup> Computer and Information Sciences  
Georgia Regents University  
Augusta, GA

<sup>2</sup> Nuance Communications  
Sunnyvale, CA

**Abstract.** Full Intuitionistic Linear Logic (FILL) was first introduced by Hyland and de Paiva, and went against current beliefs that it was not possible to incorporate all of the linear connectives, e.g. tensor, par, and implication, into an intuitionistic linear logic. It was shown that their formalization of FILL did not enjoy cut-elimination by Bierman, but Bellin proposed a change to the definition of FILL in the hope to regain cut-elimination. In this note we adopt Bellin’s proposed change and give a direct proof of cut-elimination. Then we show that a categorical model of FILL in the basic dialectica category is also a LNL model of Benton and a full tensor model of Melliès’ and Tabareau’s tensorial logic. Lastly, we give a double-negation translation of linear logic into FILL that explicitly uses par in addition to tensor.

**Keywords:** full intuitionistic linear logic, classical linear logic, dialectica category, cut-elimination, tensorial logic, linear/non-linear models, categorical model, proof theory, par

## 1 Introduction

A commonly held belief during the early history of linear logic was that the linear-connective par could not be incorporated into an intuitionistic linear logic. This belief was challenged when de Paiva gave a categorical understanding of Gödel’s Dialectica interpretation in terms of dialectica categories [9,8].

Dialectica categories were initially believed to be models of intuitionistic logic, but they are actually models of intuitionistic linear logic, containing the linear connectives: tensor, implication, the additives, and the exponentials. Further work improved de Paiva’s models to capture both intuitionistic and classical linear logic. Armed with this semantic insight de Paiva gave the first formalization of Full Intuitionistic Linear Logic (FILL) [8]. FILL is a sequent calculus with multiple conclusions in addition to multiple hypotheses. Logics of this type go back to Gentzen’s work on the sequent calculus for classical logic LK and for intuitionistic logic LJ, and Maehara’s work on LJ’ [15,23]. The sequents in these types of logics usually have the form  $\Gamma \vdash \Delta$  where  $\Gamma$  and  $\Delta$  are multisets of formulas. Sequents such as these are read as “the conjunction of the formulas in  $\Gamma$  imply the disjunction of the formulas in  $\Delta$ ”. For a brief, but more complete history of logics with multiple conclusions see the introduction to [11].

Gentzen showed that to obtain intuitionistic logic one could start with the logic LK and then place a cardinality restriction on the right-hand side of sequents, however, this is not the only means of enforcing intuitionism. Maehara showed that in the propositional case one could simply place the cardinality restriction on the premise of the implication right rule, and leave all of the other rules of LK unrestricted. This restriction is sometimes called the Dragalin restriction, as it appeared in his AMS textbook [12]. The classical implication right rule has the form:

$$\frac{\Gamma, A \vdash B, \Delta}{\Gamma \vdash A \multimap B, \Delta} \text{ IMPR}$$

By placing the Dragalin restriction on the previous rule we obtain:

$$\frac{\Gamma, A \vdash B}{\Gamma \vdash A \multimap B} \text{IMPR}$$

de Paiva’s first formalization of FILL used the Dragalin restriction, see [8] p. 58, but Schellinx showed that this restriction has the unfortunate consequence of breaking cut-elimination [21].

Later, Hyland and de Paiva gave an alternative formalization of FILL with the intention of regaining cut-elimination [13]. This new formalization lifted the Dragalin restriction by decorating sequents with a term assignment. Hypotheses were assigned variables, and the conclusions were assigned terms. Then using these terms one can track the use of hypotheses throughout a derivation. They proposed a new implication right rule:

$$\frac{\Gamma, x : A \vdash t : B, \Delta \quad x \notin \text{FV}(\Delta)}{\Gamma \vdash \lambda x.t : A \multimap B, \Delta} \text{IMPR}$$

Intuitionism is enforced in this rule by requiring that the variable being discharged,  $x$ , is not free terms annotating other conclusions. Unfortunately, this formalization did not enjoy cut-elimination either.

Bierman was able to give a counterexample to cut-elimination [4]. As Bierman explains the problem was with the left rule for the multiplicative disjunction par. The original rule was as follows:

$$\frac{\Gamma, x : A \vdash \Delta \quad \Gamma', y : B \vdash \Delta'}{\Gamma, \Gamma', z : A \wp B \vdash \text{let } z \text{ be } (x \wp -) \text{ in } \Delta \mid \text{let } z \text{ be } (- \wp y) \text{ in } \Delta'} \text{PARL}$$

In this rule the pattern variables  $x$  and  $y$  are bound in each term of  $\Delta$  and  $\Delta'$  respectively. Notice that the variable  $z$  becomes free in every term in  $\Delta$  and  $\Delta'$ . Bierman showed that this rule mixed with the restriction on implication right prevents the usual cut-elimination step that commutes cut with the left rule for par. The main idea behind the counterexample is that in the derivation before commuting the cut it is possible to discharge  $z$  using implication right, but after the cut is commuted past the left rule for par, the variable  $z$  becomes free in more than one conclusion, and thus, can no longer be discharged.

In the conclusion of Bierman’s note he gives an alternate left rule for par that he attributes to Bellin. This new left-rule is as follows:

$$\frac{\Gamma, x : A \vdash \Delta \quad \Gamma', y : B \vdash \Delta'}{\Gamma, \Gamma', z : A \wp B \vdash \text{let-pat } z (x \wp -) \Delta \mid \text{let-pat } z (- \wp y) \Delta'} \text{PARL}$$

In this rule  $\text{let-pat } z (x \wp -) t$  and  $\text{let-pat } z (- \wp y) t'$  only let-bind  $z$  in  $t$  or  $t'$  if  $x \in \text{FV}(t)$  or  $y \in \text{FV}(t')$ . Otherwise the terms are left unaltered. Bellin showed that by adopting this rule cut-elimination can be proven by reduction to the cut-elimination procedure for proof nets for multiplicative linear logic with the mix rule [1]. However, this is an indirect proof that requires the adoption of proof nets.

**Contributions.** In this paper our main contribution is to give a direct proof of cut-elimination for FILL with Bellin’s proposed par-left rule (Section 3). A direct proof accomplishes two goals: the first is to complete the picture of FILL Hyland and de Paiva started, and the second is to view a direct proof of cut-elimination as a means of checking the correctness of the formulation of FILL given here. The latter point is important for future work. Following the proof of cut-elimination we show that the categorical model of FILL called  $\text{Dial}_2(\text{Sets})$ , the basic dialectica category, is also a linear/non-linear model of Benton (Section 4) and a full tensor model of Mellies’ and Tabareau’s tensor logic (Section 5). Finally, we give a double-negation translation of multi-conclusion classical linear logic into FILL (Section 5.1). Due to the complexities of working in  $\text{Dial}_2(\text{Sets})$  we have formalized all of the constructions and proofs used in Section 4 and Section 5 – although our formal verification does not include the double-negation translation in Section 5.1 – in the Agda proof assistant<sup>3</sup>.

**Related Work.** The first formalization of FILL with cut-elimination was due to Braüner and de Paiva [5]. Their formalization can be seen as a linear version of LK with a sophisticated meta-level dependency tracking system. A proof of a FILL sequent in their formalization amounts to a classical derivation,  $\pi$ , invariant in what they call the FILL property:

<sup>3</sup> The Agda development can be found at <https://github.com/heades/cut-fill-agda>.

- The hypothesis discharged by an application of the implication right rule in  $\pi$  is a dependency of the conclusion of the implication being introduced.

They were able to show that their formalization is sound, complete, and enjoys cut-elimination. In favor of the term assignment formalization given here over Braüner and de Paiva’s formalization we can say that the dependency tracking system complicates both the definition of the logic and its use. However, one might conjecture that their system is more fundamental and hence more generalizable. It might be possible to prove cut-elimination of the term assignment formalization of FILL relative to Braüner and de Paiva’s dependency tracking system by erasing the terms on conclusions and then tracking which variable is free in which conclusion. However, as we stated above a direct proof is more desirable than a relative one.

de Paiva and Pereira used annotations on the sequents of LK to arrive at full intuitionistic logic (FIL) with multiple conclusion that enjoys cut-elimination [11]. They annotate hypothesis with natural number indices, and conclusions with finite sets of indices. The sets of indices on conclusions correspond to the collection of the hypotheses that the conclusion depends on. Then they have a similar property to that of Braüner and de Paiva’s formalization. In fact, the dependency tracking system is very similar to this formalization, but the dependency tracking has been collapsed into the object language instead of being at the meta-level.

Clouston et al. give both a deep inference calculus and a display calculus for FILL that admits cut-elimination [6]. Both of these systems are refinements of a larger one called bi-intuitionistic linear logic (BiLL). This logic contains every logical connective of FILL with the addition of the exclusion (or subtraction) connective. This connective can be defined categorically as the left-adjoint to par. Thus, exclusion is the dual to implication. A positive aspect to this work is that the resulting systems are annotation free, but at a price of complexity. Deep inference and display calculi are harder to understand, and their system requires FILL to be defined as a refinement of a system with additional connectives. We show in this paper that such a refinement is unnecessary. In addition, a term assignment system is closer to traditional logic than deep inference and display calculi, and it is closer, through the lens of the Curry-Howard-Lambek correspondence, to a type theoretic understanding of FILL.

## 2 Full Intuitionistic Linear Logic (FILL)

In this section we give a brief description of FILL. We first give the syntax of formulas, patterns, terms, and contexts. Following the syntax we define several meta-functions that will be used when defining the inference rules of the logic.

**Definition 1.** *The syntax for FILL is as follows:*

$$\begin{array}{ll}
(\text{Formulas}) & A, B, C, D, E ::= \top \mid \perp \mid A \multimap B \mid A \otimes B \mid A \wp B \\
(\text{Patterns}) & p ::= * \mid - \mid x \mid p_1 \otimes p_2 \mid p_1 \wp p_2 \\
(\text{Terms}) & t, e ::= x \mid * \mid \circ \mid t_1 \otimes t_2 \mid t_1 \wp t_2 \mid \lambda x. t \mid \text{let } t \text{ be } p \text{ in } e \mid t_1 t_2 \\
(\text{Left Contexts}) & \Gamma ::= \cdot \mid x : A \mid \Gamma_1, \Gamma_2 \\
(\text{Right Contexts}) & \Delta ::= \cdot \mid t : A \mid \Delta_1, \Delta_2
\end{array}$$

The formulas of FILL are standard, but we denote the unit of tensor as  $\top$  and the unit of par as  $\perp$ . Patterns are used to distinguish between the various let-expressions for tensor, par, and their units. There are three different let-expressions:

$$\begin{array}{lll}
\text{Tensor:} & \text{Par:} & \text{Tensor Unit:} \\
\text{let } t \text{ be } p_1 \otimes p_2 \text{ in } e & \text{let } t \text{ be } p_1 \wp p_2 \text{ in } e & \text{let } t \text{ be } * \text{ in } e
\end{array}$$

In addition, each of these will have their own equational rules, see Figure 2. The role each term plays in the overall logic will become clear after we introduce the inference rules.

At this point we introduce some syntax and meta-level functions that will be used in the definition of the inference rules for FILL. Left contexts are multisets of formulas each labeled with a variable, and right contexts are multisets of formulas each labeled with a term. We will often write  $\Delta_1 \mid \Delta_2$  as syntactic sugar

for  $\Delta_1, \Delta_2$ . The former should be read as “ $\Delta_1$  or  $\Delta_2$ .” We denote the usual capture-avoiding substitution by  $[t/x]t'$ , and its straightforward extension to right contexts as  $[t/x]\Delta$ . Similarly, we find it convenient to be able to do this style of extension for the let-binding as well.

**Definition 2.** *We extend let-binding terms to right contexts as follows:*

$$\begin{aligned} \text{let } t \text{ be } p \text{ in } \cdot &= \cdot \\ \text{let } t \text{ be } p \text{ in } (t' : A) &= (\text{let } t \text{ be } p \text{ in } t') : A \\ \text{let } t \text{ be } p \text{ in } (\Delta_1 \mid \Delta_2) &= (\text{let } t \text{ be } p \text{ in } \Delta_1) \mid (\text{let } t \text{ be } p \text{ in } \Delta_2) \end{aligned}$$

Lastly, we denote the usual function that computes the set of free variables in a term by  $\text{FV}(t)$ , and its straightforward extension to right contexts as  $\text{FV}(\Delta)$ .

The inference rules for FILL are defined in Figure 1. The  $\text{PARL}$  rule depends on the function  $\text{let-pat } z p \Delta$

$$\begin{array}{c} \frac{}{x : A \vdash x : A} \text{AX} \quad \frac{\Gamma \vdash t : A \mid \Delta \quad \Gamma', y : A \vdash \Delta'}{\Gamma, \Gamma' \vdash \Delta \mid [t/y]\Delta'} \text{CUT} \quad \frac{\Gamma \vdash \Delta}{\Gamma, x : \top \vdash \text{let } x \text{ be } * \text{ in } \Delta} \text{TL} \quad \frac{}{\cdot \vdash * : \top} \text{TR} \\ \\ \frac{\Gamma, x : A, y : B \vdash \Delta}{\Gamma, z : A \otimes B \vdash \text{let } z \text{ be } x \otimes y \text{ in } \Delta} \text{TENL} \quad \frac{\Gamma \vdash e : A \mid \Delta \quad \Gamma' \vdash f : B \mid \Delta'}{\Gamma, \Gamma' \vdash e \otimes f : A \otimes B \mid \Delta \mid \Delta'} \text{TENR} \quad \frac{}{x : \perp \vdash \cdot} \text{PL} \quad \frac{\Gamma \vdash \Delta}{\Gamma \vdash \circ : \perp \mid \Delta} \text{PR} \\ \\ \frac{\Gamma, x : A \vdash \Delta \quad \Gamma', y : B \vdash \Delta'}{\Gamma, \Gamma', z : A \wp B \vdash \text{let-pat } z (x \wp -) \Delta \mid \text{let-pat } z (- \wp y) \Delta'} \text{PARL} \quad \frac{\Gamma \vdash \Delta \mid e : A \mid f : B \mid \Delta'}{\Gamma \vdash \Delta \mid e \wp f : A \wp B \mid \Delta'} \text{PARR} \\ \\ \frac{\Gamma \vdash e : A \mid \Delta \quad \Gamma', x : B \vdash \Delta'}{\Gamma, y : A \multimap B, \Gamma' \vdash \Delta \mid [y e/x]\Delta'} \text{IMPL} \quad \frac{\Gamma, x : A \vdash e : B \mid \Delta \quad x \notin \text{FV}(\Delta)}{\Gamma \vdash \lambda x. e : A \multimap B \mid \Delta} \text{IMPR} \quad \frac{\Gamma, x : A, y : B \vdash \Delta}{\Gamma, y : B, x : A \vdash \Delta} \text{EXL} \\ \\ \frac{\Gamma \vdash \Delta_1 \mid t_1 : A \mid t_2 : B \mid \Delta_2}{\Gamma \vdash \Delta_1 \mid t_2 : B \mid t_1 : A \mid \Delta_2} \text{EXR} \end{array}$$

**Fig. 1.** Inference rules for FILL

which we define next.

**Definition 3.** *The function  $\text{let-pat } z p t$  is defined as follows:*

$$\begin{aligned} \text{let-pat } z (x \wp -) t &= t & \text{let-pat } z (- \wp y) t &= t & \text{let-pat } z p t &= \text{let } z \text{ be } p \text{ in } t \\ \text{where } x &\notin \text{FV}(t) & \text{where } y &\notin \text{FV}(t) \end{aligned}$$

It is straightforward to extend the previous definition to right-contexts, and we denote this extension by  $\text{let-pat } z p \Delta$ .

The motivation behind this function is that it only binds the pattern variables in  $x \wp -$  and  $- \wp y$  if and only if those pattern variables are free in the body of the let. This overcomes the counterexample given by Bierman in [4].

The terms of FILL are equipped with an equivalence relation defined in Figure 2. There are a number of  $\alpha$ ,  $\beta$ , and  $\eta$  like rules as well as several rules we call naturality rules. These rules are similar to the rules presented in [13].

### 3 Cut-elimination

FILL can be viewed from two different angles: i. as an intuitionistic linear logic with par, or ii. as a restricted form of classical linear logic. Thus, to prove cut-elimination of FILL one only needs to start with the cut-elimination procedure for intuitionistic linear logic, and then dualize all of the steps in the procedure for tensor and its unit to obtain the steps for par and its unit. Similarly, one could just as easily start with the

$$\begin{array}{c}
\frac{y \notin \text{FV}(t)}{t = [y/x]t} \text{ ALPHA} \quad \frac{x \notin \text{FV}(f)}{(\lambda x.f x) = f} \text{ ETAFUN} \quad \frac{}{(\lambda x.e) e' = [e'/x]e} \text{ BETA FUN} \quad \frac{}{\text{let } * \text{ be } * \text{ in } e = e} \text{ ETAI1} \\
\frac{y \notin \text{FV}(f)}{\text{let } y \text{ be } * \text{ in } f = f} \text{ ETA2I} \quad \frac{}{\text{let } u \text{ be } * \text{ in } [* / z]f = [u / z]f} \text{ BETAI} \quad \frac{}{[\text{let } u \text{ be } * \text{ in } e / y]f = \text{let } u \text{ be } * \text{ in } [e / y]f} \text{ NATI} \\
\frac{x, y \notin \text{FV}(t)}{\text{let } t' \text{ be } x \otimes y \text{ in } t = t} \text{ ETATEN} \quad \frac{}{\text{let } e \otimes t \text{ be } x \otimes y \text{ in } u = [e/x, t/y]u} \text{ BETA1TEN} \\
\frac{}{\text{let } u \text{ be } x \otimes y \text{ in } [x \otimes y / z]f = [u / z]f} \text{ BETA2TEN} \quad \frac{}{[\text{let } u \text{ be } x \otimes y \text{ in } g / w]f = \text{let } u \text{ be } x \otimes y \text{ in } [g / w]f} \text{ NATTEN} \\
\frac{}{u = \circ} \text{ ETAPARU} \quad \frac{}{(\text{let } u \text{ be } x \wp - \text{in } x) \wp (\text{let } u \text{ be } - \wp y \text{ in } y) = u} \text{ ETAPAR} \\
\frac{}{\text{let } u \wp t \text{ be } x \wp - \text{in } e = [u/x]e} \text{ BETA1PAR} \quad \frac{}{\text{let } u \wp t \text{ be } - \wp y \text{ in } e = [t/y]e} \text{ BETA2PAR} \\
\frac{}{\text{let } t \text{ be } x \wp - \text{in } [u/x]f = [\text{let } t \text{ be } x \wp - \text{in } u/x]f} \text{ NAT1PAR} \\
\frac{}{\text{let } t \text{ be } - \wp y \text{ in } [v/y]f = [\text{let } t \text{ be } - \wp y \text{ in } v/y]f} \text{ NAT2PAR} \quad \frac{t = t'}{\lambda x.t = \lambda x.t'} \text{ LAM} \quad \frac{t_1 = t'_1}{t_1 t_2 = t'_1 t_2} \text{ APP1} \\
\frac{t_2 = t'_2}{t_1 t_2 = t_1 t'_2} \text{ APP2} \quad \frac{t_1 = t'_1}{t_1 \otimes t_2 = t'_1 \otimes t_2} \text{ TEN1} \quad \frac{t_2 = t'_2}{t_1 \otimes t_2 = t_1 \otimes t'_2} \text{ TEN2} \quad \frac{t_1 = t'_1}{t_1 \wp t_2 = t'_1 \wp t_2} \text{ PAR1} \\
\frac{t_2 = t'_2}{t_1 \wp t_2 = t_1 \wp t'_2} \text{ PAR2} \quad \frac{t = t'}{\text{let } t \text{ be } p \text{ in } e = \text{let } t' \text{ be } p \text{ in } e} \text{ LET1} \quad \frac{e = e'}{\text{let } t \text{ be } p \text{ in } e = \text{let } t \text{ be } p \text{ in } e'} \text{ LET2} \\
\frac{}{t = t} \text{ REFL} \quad \frac{t = t'}{t' = t} \text{ SYM} \quad \frac{t_1 = t_2 \quad t_2 = t_3}{t_1 = t_3} \text{ TRANS}
\end{array}$$

**Fig. 2.** Equivalence on terms

cut-elimination procedure for classical linear logic, and then apply the restriction on the implication right rule producing a cut-elimination procedure for FILL.

The major difference between proving cut-elimination of FILL from classical or intuitionistic linear logic is that we must prove an invariant across each step in the procedure. The invariant is that if a derivation  $\pi$  is transformed into a derivation  $\pi'$ , then the terms in the conclusion of the final rule applied in  $\pi$  must be transformable, when the equivalences defined in Figure 2 are taken as left-to-right rewrite rules, into the terms in the conclusion of the final rule applied in  $\pi'$ .

The cut elimination procedure requires the following two basic results:

**Lemma 1 (Substitution Distribution).** *For any terms  $t$ ,  $t_1$ , and  $t_2$ ,  $[t_1/x][t_2/y]t = [[t_1/x]t_2/y][t_2/x]t$ .*

*Proof.* This proof holds by straightforward induction on the form of  $t$ .

**Lemma 2 (Let-pat Distribution).** *For any terms  $t$ ,  $t_1$ , and  $t_2$ , and pattern  $p$ ,  $\text{let-pat } t p [t_1/y]t_2 = [\text{let-pat } t p t_1/y]t_2$ .*

*Proof.* This proof holds by case splitting over  $p$ , and then using the naturality equations for the respective pattern.

We finally arrive at cut-elimination.

**Theorem 1.** *If  $\Gamma \vdash t_1 : A_1, \dots, t_i : A_i$  steps to  $\Gamma \vdash t'_1 : A_1, \dots, t'_i : A_i$  using the cut-elimination procedure, then  $t_j = t'_j$  for  $1 \leq j \leq i$ .*

*Proof.* The cut-elimination procedure given here is the standard cut-elimination procedure for classical linear logic except the cases involving the implication right rule have the FILL restriction. The structure of our procedure follows the structure of the procedure found in [16]. Throughout this proof we treat the equivalences defined in Figure 2 as left-to-right rewrite rules.

Case: commuting conversion cut vs cut (first case). The following proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \quad \frac{\frac{\pi_2}{\vdots} \quad \frac{\pi_3}{\vdots}}{\Gamma_2, x : A, \Gamma_3 \vdash t_1 : B \mid \Delta_1} \quad \frac{\Gamma_1, y : B, \Gamma_4 \vdash \Delta_2}{\Gamma_1, y : B, \Gamma_4 \vdash \Delta_2}}{\Gamma_1, \Gamma_2, x : A, \Gamma_3, \Gamma_4 \vdash \Delta_1 \mid [t_1/y]\Delta_2} \text{CUT}}{\Gamma_1, \Gamma_2, \Gamma, \Gamma_3, \Gamma_4 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x][t_1/y]\Delta_2} \text{CUT}$$

is transformed into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \quad \frac{\pi_2}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_1, y : B, \Gamma_4 \vdash \Delta_2}}{\frac{\frac{\Gamma_2, x : A, \Gamma_3 \vdash t_1 : B \mid \Delta_1}{\Gamma_2, \Gamma, \Gamma_3 \vdash [t/x]t_1 : B \mid [t/x]\Delta_1} \quad \frac{\Gamma_1, y : B, \Gamma_4 \vdash \Delta_2}{\Gamma_1, y : B, \Gamma_4 \vdash \Delta_2}}{\Gamma_1, \Gamma_2, \Gamma, \Gamma_3, \Gamma_4 \vdash \Delta \mid [t/x]\Delta_1 \mid [[t/x]t_1/y]\Delta_2} \text{CUT}} \text{CUT}$$

First, if  $\Delta_2$  is empty, then all the terms in the conclusion of the previous two derivations are equivalent. So suppose  $\Delta_2 = t_2 : C \mid \Delta'_2$ . Then we know that the term  $[t/x][t_1/y]t_2$  in the first derivation above is equivalent to  $[[t/x]t_1/y][t/x]t_2$  by Lemma 1. Furthermore, by inspecting the first derivation we can see that  $x \notin \text{FV}(t_2)$ , and thus,  $[[t/x]t_1/y][t/x]t_2 = [[t/x]t_1/y]t_2$ . This argument may be repeated for any term in  $\Delta'_2$ , and thus, we know  $[t/x][t_1/y]\Delta_2 = [[t/x]t_1/y]\Delta_2$ .

Case: commuting conversion cut vs. cut (second case). The second commuting conversion on cut begins with the proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \quad \frac{\pi_2}{\vdots} \quad \frac{\pi_3}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\Gamma' \vdash t' : B \mid \Delta'}{\Gamma_1, x : A, \Gamma_2, \Gamma' \vdash \Delta' \mid [t'/y]\Delta_1} \quad \frac{\Gamma_1, x : A, \Gamma_2, y : B, \Gamma_3 \vdash \Delta_1}{\Gamma_1, x : A, \Gamma_2, y : B, \Gamma_3 \vdash \Delta_1}}{\Gamma_1, \Gamma, \Gamma_2, \Gamma', \Gamma_3 \vdash \Delta \mid [t/x]\Delta' \mid [t/x][t'/y]\Delta_1} \text{CUT}} \text{CUT}$$

is transformed into the following proof:

$$\frac{\frac{\frac{\pi_2}{\vdots} \quad \frac{\pi_1}{\vdots} \quad \frac{\pi_3}{\vdots}}{\Gamma' \vdash t' : B \mid \Delta'} \quad \frac{\frac{\Gamma \vdash t : A \mid \Delta} \quad \frac{\Gamma_1, x : A, \Gamma_2, y : B, \Gamma_3 \vdash \Delta_1}{\Gamma_1, \Gamma, \Gamma_2, y : B, \Gamma_3 \vdash \Delta \mid [t/x]\Delta_1}}{\frac{\Gamma_1, \Gamma, \Gamma_2, \Gamma', \Gamma_3 \vdash \Delta' \mid [t'/y]\Delta \mid [t'/y][t/x]\Delta_1} \quad \frac{\Gamma_1, \Gamma, \Gamma_2, \Gamma', \Gamma_3 \vdash \Delta' \mid [t'/y]\Delta \mid [t'/y][t/x]\Delta_1}}{\Gamma_1, \Gamma, \Gamma_2, \Gamma', \Gamma_3 \vdash [t'/y]\Delta \mid \Delta' \mid [t'/y][t/x]\Delta_1} \text{CUT}} \text{CUT}$$

SERIES OF EXCHANGES

We know  $x, y \notin \text{FV}(\Delta)$  by inspection of the first derivation, and so we know that  $\Delta = [t'/y]\Delta$  and  $\Delta' = [t/x]\Delta'$ . Without loss of generality suppose  $\Delta_1 = t_1 : C \mid \Delta'_1$ . Then we know that  $x, y \notin \text{FV}(t)$  and  $x, y \notin \text{FV}(t')$ . Thus, by this fact and Lemma 1, we know that  $[t/x][t'/y]t_1 = [[t/x]t'/y][t/x]t_1 = [t'/y][t/x]t_1$ . This argument can be repeated for any term in  $\Delta'_1$ , hence,  $[t/x][t'/y]\Delta_1 = [t'/y][t/x]\Delta_1$ .

Case: the  $\eta$ -expansion cases: tensor. The proof

$$\frac{}{x : A \otimes B \vdash x : A \otimes B} \text{Ax}$$

is transformed into the proof

$$\frac{\frac{\frac{}{y : A \vdash y : A} \text{Ax} \quad \frac{}{z : B \vdash z : B} \text{Ax}}{y : A, z : B \vdash y \otimes z : A \otimes B} \text{TR}}{x : A \otimes B \vdash \text{let } x \text{ be } y \otimes z \text{ in } (y \otimes z) : A \otimes B} \text{TL}}$$

By the rule  $\text{EQ\_ETA\_TENSOR}$  we know  $\text{let } x \text{ be } y \otimes z \text{ in } (y \otimes z) = x$ .

Case: the  $\eta$ -expansion cases: par. The proof

$$\frac{}{x : A \wp B \vdash x : A \wp B} \text{Ax}$$

is transformed into the proof

$$\frac{\frac{\frac{}{y : A \vdash y : A} \text{Ax} \quad \frac{}{z : B \vdash z : B} \text{Ax}}{x : A \wp B \vdash \text{let } x \text{ be } (y \wp -) \text{ in } y : A \mid \text{let } x \text{ be } (- \wp z) \text{ in } z : B} \text{PARL}}{x : A \wp B \vdash (\text{let } x \text{ be } (y \wp -) \text{ in } y) \wp (\text{let } x \text{ be } (- \wp z) \text{ in } z) : A \wp B} \text{PARR}}$$

By rule  $\text{EQ\_ETA\_PAR}$  we know  $((\text{let } x \text{ be } (y \wp -) \text{ in } y) \wp (\text{let } x \text{ be } (- \wp z) \text{ in } z)) = x$ .

Case: the  $\eta$ -expansion cases: implication. The proof

$$\frac{}{x : A \multimap B \vdash x : A \multimap B} \text{Ax}$$

transforms into the proof

$$\frac{\frac{\frac{}{y : A \vdash y : A} \text{Ax} \quad \frac{}{z : B \vdash z : B} \text{Ax}}{y : A, x : A \multimap B \vdash x y : B} \text{IMPL}}{x : A \multimap B \vdash \lambda y. x y : A \multimap B} \text{IMPR}}$$

All terms in the two derivations are equivalent, because  $(\lambda y. x y) = x$  by the  $\text{EQ\_ETA\_FUN}$  rule.

Case: the  $\eta$ -expansion cases: tensor unit. The proof

$$\frac{}{x : \top \vdash x : \top} \text{Ax}$$

transforms into the proof

$$\frac{\frac{}{\cdot \vdash * : \top} \text{IR}}{x : \top \vdash \text{let } x \text{ be } * \text{ in } * : \top} \text{IL}}$$

We know  $x = \text{let } x \text{ be } * \text{ in } *$  by  $\text{EQ\_ETA}$ .

Case: the  $\eta$ -expansion cases: par unit. The proof

$$\frac{}{x : \perp \vdash x : \perp} \text{Ax}$$

transforms into the proof

$$\frac{\overline{x : \perp \vdash \cdot} \text{ PL}}{x : \perp \vdash \circ : \perp} \text{ PR}$$

We know  $x = \circ$  by EQ-ETAPARU.

Case: the axiom steps: the axiom step. The proof

$$\frac{\frac{\overline{x : A \vdash x : A} \text{ AX} \quad \frac{\pi \quad \vdots}{\Gamma_1, y : A, \Gamma_2 \vdash \Delta}}{\Gamma_1, x : A, \Gamma_2 \vdash [x/y]\Delta} \text{ CUT}}$$

transforms into the proof

$$\frac{\pi \quad \vdots}{\Gamma_1, y : A, \Gamma_2 \vdash \Delta}$$

By EQ-ALPHA, we know, for any  $t$  in  $\Delta$ ,  $t = [x/y]t$ , and hence  $\Delta = [x/y]\Delta$ .

Case: the axiom steps: conclusion vs. axiom. The proof

$$\frac{\frac{\pi \quad \vdots}{\Gamma \vdash t : A \mid \Delta} \quad \overline{x : A \vdash x : A} \text{ AX}}{\Gamma \vdash \Delta \mid [t/x]x : A} \text{ CUT}$$

transforms into

$$\frac{\pi \quad \vdots}{\Gamma \vdash t : A \mid \Delta} \text{ SERIES OF EXCHANGES} \\ \Gamma \vdash \Delta \mid t : A$$

By the definition of the substitution function we know  $t = [t/x]x$ .

Case: the exchange steps: conclusion vs. left-exchange (the first case). The proof

$$\frac{\frac{\pi_1 \quad \vdots}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2 \quad \vdots}{\Gamma_1, x : A, y : B, \Gamma_2 \vdash \Delta'}{\Gamma_1, y : B, x : A, \Gamma_2 \vdash \Delta'} \text{ EXL}}{\Gamma_1, y : B, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta'} \text{ CUT}}$$

transforms into the proof

$$\frac{\frac{\pi_1 \quad \vdots}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\pi_2 \quad \vdots}{\Gamma_1, x : A, y : B, \Gamma_2 \vdash \Delta'}}{\Gamma_1, \Gamma, y : B, \Gamma_2 \vdash \Delta \mid [t/x]\Delta'} \text{ CUT} \\ \Gamma_1, y : B, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta' \text{ SERIES OF EXCHANGES}$$



Clearly, all terms are equivalent.

Case: the exchange steps: conclusion vs. left-exchange (the second case). The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : B \mid \Delta} \quad \frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, y : B, \Gamma_2 \vdash \Delta'}{\Gamma_1, y : B, x : A, \Gamma_2 \vdash \Delta'} \text{EXL}}{\Gamma_1, \Gamma, x : A, \Gamma_2 \vdash \Delta \mid [t/y]\Delta'} \text{CUT}}{\Gamma_1, \Gamma, x : A, \Gamma_2 \vdash \Delta \mid [t/y]\Delta'}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : B \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, y : B, \Gamma_2 \vdash \Delta'}{\Gamma_1, x : A, \Gamma, \Gamma_2 \vdash \Delta \mid [t/y]\Delta'} \text{CUT}}{\Gamma_1, \Gamma, x : A, \Gamma_2 \vdash \Delta \mid [t/y]\Delta'} \text{SERIES OF EXCHANGES}$$

Clearly, all terms are equivalent.

Case: the exchange steps: conclusion vs. right-exchange The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2 \vdash \Delta_1 \mid t_1 : B \mid t_2 : C \mid \Delta'}{\Gamma_1, x : A, \Gamma_2 \vdash \Delta_1 \mid t_2 : C \mid t_1 : B \mid \Delta'} \text{EXR}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x]t_2 : C \mid [t/x]t_1 : B \mid [t/x]\Delta'} \text{CUT}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x]t_2 : C \mid [t/x]t_1 : B \mid [t/x]\Delta'}$$

transforms into this proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2 \vdash \Delta_1 \mid t_1 : B \mid t_2 : C \mid \Delta'}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x]t_1 : B \mid [t/x]t_2 : C \mid [t/x]\Delta'} \text{CUT}}{\Gamma_1, \Gamma, \Gamma_2 \vdash [t/x]\Delta_1 \mid [t/x]t_2 : C \mid [t/x]t_1 : B \mid [t/x]\Delta'} \text{EXR}$$

Clearly, all terms are equivalent.

Case: principal formula vs. principal formula: tensor. The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma_1 \vdash t_1 : A \mid \Delta_1} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_2 \vdash t_2 : B \mid \Delta_2} \text{TR} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, x : A, y : B, \Gamma_4 \vdash \Delta_3} \text{TL}}{\Gamma_3, \Gamma_1, \Gamma_2, \Gamma_4 \vdash \Delta_1 \mid \Delta_2 \mid [t_1 \otimes t_2/z](\text{let } z \text{ be } x \otimes y \text{ in } \Delta_3)} \text{CUT}}$$

is transformed into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma_1 \vdash t_1 : A \mid \Delta_1} \quad \frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_2 \vdash t_2 : B \mid \Delta_2} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, x : A, y : B, \Gamma_4 \vdash \Delta_3}}{\Gamma_3, x : A, \Gamma_2, \Gamma_4 \vdash \Delta_2 \mid [t_2/y]\Delta_3} \text{CUT}}{\Gamma_3, \Gamma_1, \Gamma_2, \Gamma_4 \vdash \Delta_1 \mid \Delta_2 \mid [t_1/x][t_2/y]\Delta_3} \text{CUT}$$

Without loss of generality suppose  $\Delta_3 = t_3 : C, \Delta'_3$ . We can see that  $[t_1 \otimes t_2/z](\text{let } z \text{ be } x \otimes y \text{ in } t_3) = \text{let } t_1 \otimes t_2 \text{ be } x \otimes y \text{ in } t_3$  by the definition of substitution, and by using the  $\text{EQ\_BETA1TENSOR}$  rule we obtain  $\text{let } t_1 \otimes t_2 \text{ be } x \otimes y \text{ in } t_3 = [t_1/x][t_2/y]t_3$ . This argument can be repeated for any term in  $[t_1 \otimes t_2/z](\text{let } z \text{ be } x \otimes y \text{ in } \Delta'_3)$ , and thus,  $[t_1 \otimes t_2/z](\text{let } z \text{ be } x \otimes y \text{ in } \Delta_3) = [t_1/x][t_2/y]\Delta_3$ .

Note that in the second derivation of the above transformation we first cut on  $B$ , and then  $A$ , but we could have cut on  $A$  first, and then  $B$ , but this would yield equivalent derivations as above by using Lemma 1.

Case: principal formula vs. principal formula: par. The proof

$$\text{PARR} \frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma_1 \vdash \Delta_1 \mid t_1 : A \mid t_2 : B \mid \Delta_2} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_2, x : A \vdash \Delta_3} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, y : B \vdash \Delta_4}}{\Gamma_1 \vdash \Delta_1 \mid t_1 \wp t_2 : A \wp B \mid \Delta_2} \quad \frac{\Gamma_2, \Gamma_3, z : A \wp B \vdash \text{let-pat } z (x \wp -) \Delta_3 \mid \text{let-pat } z (- \wp y) \Delta_4}{\Gamma_2, \Gamma_3, \Gamma_1 \vdash \Delta_1 \mid \Delta_2 \mid [t_1 \wp t_2/z](\text{let-pat } z (x \wp -) \Delta_3) \mid [t_1 \wp t_2/z](\text{let-pat } z (- \wp y) \Delta_4)}}{\text{CUT}} \text{PARR}$$

is transformed into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma_1 \vdash \Delta_1 \mid t_1 : A \mid t_2 : B \mid \Delta_2} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, y : B \vdash \Delta_4}}{\Gamma_3, \Gamma_1 \vdash \Delta_1 \mid t_1 : A \mid \Delta_2 \mid [t_2/y]\Delta_4} \text{CUT} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_2, x : A \vdash \Delta_3}}{\Gamma_2, \Gamma_3, \Gamma_1 \vdash \Delta_1 \mid \Delta_2 \mid [t_2/y]\Delta_4 \mid [t_1/x]\Delta_3} \text{CUT}}{\Gamma_2, \Gamma_3, \Gamma_1 \vdash \Delta_1 \mid \Delta_2 \mid [t_1/x]\Delta_3 \mid [t_2/y]\Delta_4} \text{SERIES OF EXCHANGES}$$

Without loss of generality consider the case when  $\Delta_3 = t_3 : C_1 \mid \Delta'_3$  and  $\Delta_4 = t_4 : C_2 \mid \Delta'_4$ . First,  $[t_1 \wp t_2/z](\text{let-pat } z (x \wp -) t_3) = \text{let-pat } (t_1 \wp t_2) (x \wp -) t_3$ , and by  $\text{EQ\_BETA1PAR}$  we know  $\text{let-pat } (t_1 \wp t_2) (x \wp -) t_3 = [t_1/x]t_3$  if  $x \in \text{FV}(t_3)$  or  $\text{let-pat } (t_1 \wp t_2) (x \wp -) t_3 = t_3$  otherwise. In the latter case we can see that  $t_3 = [t_1/x]t_3$ , thus, in both cases  $\text{let-pat } (t_1 \wp t_2) (x \wp -) t_3 = [t_1/x]t_3$ . This argument can be repeated for any terms in  $\Delta'_3$ , and hence  $[t_1 \wp t_2/z](\text{let-pat } z (x \wp -) \Delta_3) = \text{let-pat } (t_1 \wp t_2) (x \wp -) \Delta_3 = [t_1/x]\Delta_3$ .

We can apply a similar argument for

$[t_1 \wp t_2/z](\text{let-pat } z (- \wp y) t_4)$  and  $[t_1 \wp t_2/z](\text{let-pat } z (- \wp y) \Delta_4)$ .

Note that we could have first cut on  $A$ , and then on  $B$  in the second derivation, but we would have arrived at the same result just with potentially more exchanges on the right.

Case: principal formula vs. principal formula: implication. The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma, x : A \vdash t : B \mid \Delta} \quad \frac{x \notin \text{FV}(\Delta)}{\Gamma \vdash \lambda x.t : A \multimap B \mid \Delta} \text{IMPR} \quad \frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1 \vdash t_1 : A \mid \Delta_1} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_2, y : B \vdash \Delta_2}}{\Gamma_1, z : A \multimap B, \Gamma_2 \vdash \Delta_1 \mid [z t_1/y]\Delta_2} \text{IMPL}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [\lambda x.t/z]\Delta_1 \mid [\lambda x.t/z][z t_1/y]\Delta_2} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1 \vdash t_1 : A \mid \Delta_1} \quad \frac{\frac{\pi_1}{\vdots}}{\Gamma, x : A \vdash t : B \mid \Delta} \quad \frac{x \notin \text{FV}(\Delta)}{\Gamma, \Gamma_1 \vdash \Delta_1 \mid [t_1/x]t : B \mid [t_1/x]\Delta} \text{CUT} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_2, y : B \vdash \Delta_2}}{\Gamma_2, \Gamma, \Gamma_1 \vdash \Delta_1 \mid [t_1/x]\Delta \mid [[t_1/x]t/y]\Delta_2} \text{CUT}}{\Gamma_1, \Gamma, \Gamma_2 \vdash [t_1/x]\Delta \mid \Delta_1 \mid [[t_1/x]t/y]\Delta_2} \text{SERIES OF EXCHANGES}$$

Without loss of generality consider the case when  $\Delta_2 = t_2 : C \mid \Delta'_2$ . First, by hypothesis we know  $x \notin \text{FV}(\Delta)$ , and so we know  $\Delta = [t_1/x]\Delta$ . We can see that  $[\lambda x.t/z][z t_1/y]t_2 = [(\lambda x.t) t_1/y]t_2 = [[t_1/x]t/y]t_2$  by using the congruence rules of equality and the rule  $\text{EQ\_BETA1FUN}$ . This argument can be repeated for any term in  $[\lambda x.t/z][z t_1/y]\Delta'_2$ , and so  $[\lambda x.t/z][z t_1/y]\Delta_2 = [[t_1/x]t/y]\Delta_2$ . Finally, by inspecting the previous derivations we can see that  $z \notin \text{FV}(\Delta_1)$ , and thus,  $\Delta_1 = [\lambda x.t/z]\Delta_1$ .

Case: principal formula vs. principal formula: tensor unit. The proof

$$\frac{\frac{\cdot \vdash * : \top}{\cdot \vdash * : \top} \text{IR} \quad \frac{\frac{\pi}{\vdots} \overline{\Gamma \vdash \Delta}}{\Gamma, x : \top \vdash \text{let } x \text{ be } * \text{ in } \Delta} \text{IL}}{\Gamma \vdash [* / x](\text{let } x \text{ be } * \text{ in } \Delta)} \text{CUT}$$

is transformed into the proof

$$\frac{\pi}{\vdots} \overline{\Gamma \vdash \Delta}$$

Without loss of generality suppose  $\Delta = t : A \mid \Delta'$ . We can see that  $[* / x](\text{let } x \text{ be } * \text{ in } t) = \text{let } * \text{ be } * \text{ in } t = t$  by the definition of substitution and the  $\text{EQ}_{\text{ETA}}$  rule. This argument can be repeated for any term in  $[* / x](\text{let } x \text{ be } * \text{ in } \Delta')$ , and hence,  $[* / x](\text{let } x \text{ be } * \text{ in } \Delta) = \Delta$ .

Case: principal formula vs. principal formula: par unit. The proof

$$\frac{\frac{\frac{\pi}{\vdots} \overline{\Gamma \vdash \Delta}}{\Gamma \vdash \circ : \perp \mid \Delta} \text{PR} \quad \frac{\overline{x : \perp \vdash \cdot}}{x : \perp \vdash \cdot} \text{PL}}{\Gamma \vdash \Delta \mid [\circ / x] \cdot} \text{CUT}$$

transforms into the proof

$$\frac{\pi}{\vdots} \overline{\Gamma \vdash \Delta}$$

Clearly,  $[\circ / x] \cdot = \cdot$ .

Case: secondary conclusion: left introduction of implication. The proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \overline{\Gamma \vdash t_1 : A \mid \Delta}}{\Gamma, y : A \multimap B, \Gamma_1, \Gamma_2 \vdash \Delta \mid [y t_1 / x] t_2 : C \mid [y t_1 / x] \Delta_2} \text{IMPL} \quad \frac{\frac{\pi_2}{\vdots} \overline{\Gamma_1, x : B, \Gamma_2 \vdash t_2 : C \mid \Delta_2}}{\Gamma_3, z : C, \Gamma_4 \vdash \Delta_3} \text{CUT}}{\Gamma_3, \Gamma, y : A \multimap B, \Gamma_1, \Gamma_2, \Gamma_4 \vdash \Delta \mid [y t_1 / x] \Delta_2 \mid [[y t_1 / x] t_2 / z] \Delta_3} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \overline{\Gamma \vdash t_1 : A \mid \Delta}}{\Gamma, y : A \multimap B, \Gamma_3, \Gamma_1, \Gamma_2, \Gamma_4 \vdash \Delta \mid [y t_1 / x] \Delta_2 \mid [y t_1 / x][t_2 / z] \Delta_3} \text{IMPL} \quad \frac{\frac{\frac{\pi_2}{\vdots} \overline{\Gamma_1, x : B, \Gamma_2 \vdash t_2 : C \mid \Delta_2}}{\Gamma_3, \Gamma_1, x : B, \Gamma_2, \Gamma_4 \vdash \Delta_2 \mid [t_2 / z] \Delta_3} \text{CUT} \quad \frac{\frac{\pi_3}{\vdots} \overline{\Gamma_3, z : C, \Gamma_4 \vdash \Delta_3}}{\Gamma_3, \Gamma, y : A \multimap B, \Gamma_1, \Gamma_2, \Gamma_4 \vdash \Delta \mid [y t_1 / x] \Delta_2 \mid [y t_1 / x][t_2 / z] \Delta_3} \text{SERIES OF EXCHANGES}}{\Gamma_3, \Gamma, y : A \multimap B, \Gamma_1, \Gamma_2, \Gamma_4 \vdash \Delta \mid [y t_1 / x] \Delta_2 \mid [y t_1 / x][t_2 / z] \Delta_3} \text{SERIES OF EXCHANGES}$$

This case holds because we can prove that  $[y t_1 / x][t_2 / z] \Delta_3 = [[y t_1 / x] t_2 / z] \Delta_3$  by Lemma 1 and the fact that  $x \notin \text{FV}(\Delta_3)$ .

Case: secondary conclusion: left introduction of exchange. The proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \quad \frac{\Gamma, y : B, x : A, \Gamma' \vdash t : C \mid \Delta}{\Gamma, x : A, y : B, \Gamma' \vdash t : C \mid \Delta} \text{EXL}}{\Gamma_1, \Gamma, x : A, y : B, \Gamma', \Gamma_2 \vdash \Delta \mid [t/z]\Delta_2} \text{CUT}}{\Gamma_1, z : C, \Gamma_2 \vdash \Delta_2} \pi_2$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \quad \frac{\Gamma, y : B, x : A, \Gamma' \vdash t : C \mid \Delta}{\Gamma_1, \Gamma, y : B, x : A, \Gamma', \Gamma_2 \vdash \Delta \mid [t/z]\Delta_2} \text{CUT}}{\Gamma_1, \Gamma, x : A, y : B, \Gamma', \Gamma_2 \vdash \Delta \mid [t/z]\Delta_2} \text{EXL}}{\Gamma_1, z : C, \Gamma_2 \vdash \Delta_2} \pi_2$$

Clearly, all terms are equivalent.

Case: secondary conclusion: left introduction of tensor. The proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \quad \frac{\Gamma, x : A, y : B \vdash t : C \mid \Delta}{\Gamma, z : A \otimes B \vdash \text{let } z \text{ be } x \otimes y \text{ in } t : C \mid \text{let } z \text{ be } x \otimes y \text{ in } \Delta} \text{TL}}{\Gamma_1, \Gamma, z : A \otimes B, \Gamma_2 \vdash \text{let } z \text{ be } x \otimes y \text{ in } \Delta \mid [\text{let } z \text{ be } x \otimes y \text{ in } t/w]\Delta_2} \text{CUT}}{\Gamma_1, w : C, \Gamma_2 \vdash \Delta_2} \pi_2$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \quad \frac{\Gamma, x : A, y : B \vdash t : C \mid \Delta}{\Gamma_1, \Gamma, x : A, y : B, \Gamma_2 \vdash \Delta \mid [t/w]\Delta_2} \text{CUT}}{\Gamma_1, \Gamma, z : A \otimes B, \Gamma_2 \vdash \text{let } z \text{ be } x \otimes y \text{ in } \Delta \mid \text{let } z \text{ be } x \otimes y \text{ in } ([t/w]\Delta_2)} \text{TL}}{\Gamma_1, w : C, \Gamma_2 \vdash \Delta_2} \pi_2$$

It suffices to show that  $\text{let } z \text{ be } x \otimes y \text{ in } ([t/w]\Delta_2) = [\text{let } z \text{ be } x \otimes y \text{ in } t/w]\Delta_2$ . This is a simple consequence of the rule  $\text{EQ\_NAT\_TENSOR}$ .

Case: secondary conclusion: left introduction of Par The proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \quad \frac{\Gamma', y : B \vdash t' : C \mid \Delta'}{\Gamma, \Gamma', z : A \wp B \vdash \text{let-pat } z (x \wp -) \Delta \mid \text{let-pat } z (- \wp y) t' : C \mid \text{let-pat } z (- \wp y) \Delta'} \text{PARL}}{\Gamma_1, \Gamma, \Gamma', z : A \wp B, \Gamma_2 \vdash \text{let-pat } z (x \wp -) \Delta \mid \text{let-pat } z (- \wp y) \Delta' \mid [\text{let-pat } z (- \wp y) t'/w]\Delta_2} \text{CUT}}{\Gamma_1, w : C, \Gamma_2 \vdash \Delta_2} \pi_3$$

is transformed into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \quad \frac{\frac{\pi_2}{\vdots} \quad \frac{\Gamma', y : B \vdash t' : C \mid \Delta'}{\Gamma_1, \Gamma', y : B, \Gamma_2 \vdash \Delta' \mid [t'/w]\Delta_2} \text{CUT}}{\Gamma, \Gamma_1, \Gamma', \Gamma_2, z : A \wp B \vdash \text{let-pat } z (x \wp -) \Delta \mid \text{let-pat } z (- \wp y) \Delta' \mid \text{let-pat } z (- \wp y) [t'/w]\Delta_2} \text{PARL}}{\Gamma_1, \Gamma, \Gamma', z : A \wp B, \Gamma_2 \vdash \text{let-pat } z (x \wp -) \Delta \mid \text{let-pat } z (- \wp y) \Delta' \mid \text{let-pat } z (- \wp y) [t'/w]\Delta_2} \text{SERIES OF EXCHANGES}}$$

It suffices to show that  $\text{let-pat } z(-\wp y)[t'/w]\Delta_2 = [\text{let-pat } z(-\wp y)t'/w]\Delta_2$ . This follows from the rule  $\text{EQ\_NAT2PAR}$ .

Case: secondary conclusion: left introduction of tensor unit. The proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \overline{\Gamma \vdash t : C \mid \Delta}}{\Gamma, x : \top \vdash t : C \mid \Delta} \text{IL} \quad \frac{\pi_2}{\vdots} \overline{\Gamma_1, w : C, \Gamma_2 \vdash \Delta_1}}{\Gamma_1, \Gamma, x : \top, \Gamma_2 \vdash \Delta \mid [t/w]\Delta_1} \text{CUT}$$

is transformed into the following:

$$\frac{\frac{\frac{\pi_1}{\vdots} \overline{\Gamma \vdash t : C \mid \Delta} \quad \frac{\pi_2}{\vdots} \overline{\Gamma_1, w : C, \Gamma_2 \vdash \Delta_1}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/w]\Delta_1} \text{CUT}}{\frac{\Gamma_1, \Gamma, \Gamma_2, x : \top \vdash \Delta \mid [t/w]\Delta_1}{\Gamma_1, \Gamma, x : \top, \Gamma_2 \vdash \Delta \mid [t/w]\Delta_1} \text{IL}} \text{SERIES OF EXCHANGES}$$

Clearly, all terms are equivalent. Note that we do not give a case for secondary conclusion of the left introduction of par's unit, because it can only be introduced given an empty right context, and thus there is no cut formula.

Case: secondary hypothesis: left introduction of tensor. The proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \overline{\Gamma \vdash t : A \mid \Delta} \quad \frac{\pi_2}{\vdots} \overline{\Gamma_1, x : A, \Gamma_2, y : B, z : C, \Gamma_3 \vdash t_1 : D \mid \Delta_1}}{\Gamma_1, x : A, \Gamma_2, w : B \otimes C, \Gamma_3 \vdash \text{let } w \text{ be } y \otimes z \text{ in } t_1 : D \mid \text{let } w \text{ be } y \otimes z \text{ in } \Delta_1} \text{TL}}{\Gamma_1, \Gamma, \Gamma_2, w : B \otimes C, \Gamma_3 \vdash \Delta \mid [t/x](\text{let } w \text{ be } y \otimes z \text{ in } t_1) : D \mid [t/x](\text{let } w \text{ be } y \otimes z \text{ in } \Delta_1)} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \overline{\Gamma \vdash t : A \mid \Delta} \quad \frac{\pi_2}{\vdots} \overline{\Gamma_1, x : A, \Gamma_2, y : B, z : C, \Gamma_3 \vdash t_1 : D \mid \Delta_1}}{\Gamma_1, \Gamma, \Gamma_2, y : B, z : C, \Gamma_3 \vdash \Delta \mid [t/x]t_1 : D \mid [t/x]\Delta_1} \text{CUT}}{\Gamma_1, \Gamma, \Gamma_2, w : B \otimes C, \Gamma_3 \vdash \text{let } w \text{ be } x \otimes y \text{ in } \Delta \mid \text{let } w \text{ be } x \otimes y \text{ in } [t/x]t_1 : D \mid \text{let } w \text{ be } x \otimes y \text{ in } [t/x]\Delta_1} \text{TL}$$

First, we can see by inspection of the previous derivations that  $x, y \notin \text{FV}(\Delta)$ , thus, by using similar reasoning as above we can use the  $\text{ETA\_TENSOR}$  rule to obtain  $\text{let } w \text{ be } x \otimes y \text{ in } \Delta = \Delta$ . It is a well-known property of substitution that  $[t/x](\text{let } w \text{ be } x \otimes y \text{ in } t_1) = \text{let } [t/x]w \text{ be } x \otimes y \text{ in } [t/x]t_1 = \text{let } w \text{ be } x \otimes y \text{ in } [t/x]t_1$ .

Case: secondary hypothesis: right introduction of tensor (first case). The proof

$$\frac{\frac{\frac{\pi_1}{\vdots} \overline{\Gamma \vdash t : A \mid \Delta} \quad \frac{\pi_2}{\vdots} \overline{\Gamma_1, x : A, \Gamma_2 \vdash t_1 : B \mid \Delta_1} \quad \frac{\pi_3}{\vdots} \overline{\Gamma_3 \vdash t_2 : C \mid \Delta_2}}{\Gamma_1, x : A, \Gamma_2, \Gamma_3 \vdash t_1 \otimes t_2 : B \otimes C \mid \Delta_1 \mid \Delta_2} \text{TR}}{\Gamma_1, \Gamma, \Gamma_2, \Gamma_3 \vdash \Delta \mid [t/x](t_1 \otimes t_2) : B \otimes C \mid [t/x]\Delta_1 \mid [t/x]\Delta_2} \text{CUT}$$

transforms into the proof

$$\begin{array}{c}
\pi_1 \\
\vdots \\
\hline
\Gamma \vdash t : A \mid \Delta \\
\hline
\pi_2 \\
\vdots \\
\hline
\Gamma_1, x : A, \Gamma_2 \vdash t_1 : B \mid \Delta_1 \\
\hline
\pi_3 \\
\vdots \\
\hline
\Gamma_3 \vdash t_2 : C \mid \Delta_2 \\
\hline
\text{CUT} \\
\hline
\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]t_1 : B \mid [t/x]\Delta_1 \\
\hline
\text{TR} \\
\hline
\Gamma_1, \Gamma, \Gamma_2, \Gamma_3 \vdash [t/x]t_1 \otimes t_2 : B \otimes C \mid \Delta \mid [t/x]\Delta_1 \mid \Delta_2 \\
\hline
\Gamma_1, \Gamma, \Gamma_2, \Gamma_3 \vdash \Delta \mid ([t/x]t_1) \otimes t_2 : B \otimes C \mid [t/x]\Delta_1 \mid \Delta_2 \\
\hline
\text{SERIES OF EXCHANGES}
\end{array}$$

By inspection of the previous derivations we can see that  $x \notin \text{FV}(t_2)$  and  $x \notin \text{FV}(\Delta_2)$ . Thus,  $[t/x]\Delta_2 = \Delta_2$  and  $[t/x](t_1 \otimes t_2) = ([t/x]t_1) \otimes ([t/x]t_2) = ([t/x]t_1) \otimes t_2$ .

Case: secondary hypothesis: right introduction of tensor (second case). The proof

$$\begin{array}{c}
\pi_1 \\
\vdots \\
\hline
\Gamma \vdash t : A \mid \Delta \\
\hline
\pi_2 \\
\vdots \\
\hline
\Gamma_1 \vdash t_1 : B \mid \Delta_1 \\
\hline
\pi_3 \\
\vdots \\
\hline
\Gamma_2, x : A, \Gamma_3 \vdash t_2 : C \mid \Delta_2 \\
\hline
\text{TR} \\
\hline
\Gamma_1, \Gamma_2, x : A, \Gamma_3 \vdash t_1 \otimes t_2 : B \otimes C \mid \Delta_1 \mid \Delta_2 \\
\hline
\text{CUT} \\
\hline
\Gamma_1, \Gamma, \Gamma_2, \Gamma_3 \vdash \Delta \mid [t/x](t_1 \otimes t_2) : B \otimes C \mid [t/x]\Delta_1 \mid [t/x]\Delta_2
\end{array}$$

transforms into the proof

$$\begin{array}{c}
\pi_2 \\
\vdots \\
\hline
\Gamma \vdash t_1 : B \mid \Delta_1 \\
\hline
\pi_1 \\
\vdots \\
\hline
\Gamma \vdash t : A \mid \Delta \\
\hline
\pi_3 \\
\vdots \\
\hline
\Gamma_2, x : A, \Gamma_3 \vdash t_2 : C \mid \Delta_2 \\
\hline
\text{CUT} \\
\hline
\Gamma_2, \Gamma, \Gamma_3 \vdash \Delta \mid [t/x]t_2 : C \mid [t/x]\Delta_2 \\
\hline
\text{TR} \\
\hline
\Gamma_1, \Gamma_2, \Gamma, \Gamma_3 \vdash t_1 \otimes ([t/x]t_2) : B \otimes C \mid \Delta_1 \mid \Delta \mid [t/x]\Delta_2 \\
\hline
\Gamma_1, \Gamma, \Gamma_2, \Gamma_3 \vdash \Delta \mid t_1 \otimes ([t/x]t_2) : B \otimes C \mid \Delta_1 \mid [t/x]\Delta_2 \\
\hline
\text{SERIES OF EXCHANGES}
\end{array}$$

This case is similar to the previous case.

Case: secondary hypothesis: right introduction of par. The proof

$$\begin{array}{c}
\pi_1 \\
\vdots \\
\hline
\Gamma \vdash t : A \mid \Delta \\
\hline
\pi_2 \\
\vdots \\
\hline
\Gamma_1, x : A, \Gamma_2 \vdash \Delta_1 \mid t_1 : B \mid t_2 : C \mid \Delta_2 \\
\hline
\text{PARR} \\
\hline
\Gamma_1, x : A, \Gamma_2 \vdash \Delta_1 \mid t_1 \wp t_2 : B \wp C \mid \Delta_2 \\
\hline
\text{CUT} \\
\hline
\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x](t_1 \wp t_2) : B \wp C \mid [t/x]\Delta_2
\end{array}$$

transforms into the proof

$$\begin{array}{c}
\pi_1 \\
\vdots \\
\hline
\Gamma \vdash t : A \mid \Delta \\
\hline
\pi_2 \\
\vdots \\
\hline
\Gamma_1, x : A, \Gamma_2 \vdash \Delta_1 \mid t_1 : B \mid t_2 : C \mid \Delta_2 \\
\hline
\text{CUT} \\
\hline
\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x]t_1 : B \mid [t/x]t_2 : C \mid [t/x]\Delta_2 \\
\hline
\text{PARR} \\
\hline
\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x]t_1 \wp [t/x]t_2 : B \wp C \mid [t/x]\Delta_2
\end{array}$$

Clearly,  $[t/x](t_1 \wp t_2) = ([t/x]t_1) \wp [t/x]t_2$ .

Case: secondary hypothesis: left introduction of par (first case). The proof

$$\begin{array}{c}
\pi_1 \\
\vdots \\
\hline
\Gamma \vdash t : A \mid \Delta \\
\hline
\pi_2 \\
\vdots \\
\hline
\Gamma_1, x : A, \Gamma_2, y : B \vdash \Delta_1 \\
\hline
\pi_3 \\
\vdots \\
\hline
\Gamma_3, z : C \vdash \Delta_2 \\
\hline
\text{PARR} \\
\hline
\Gamma_1, x : A, \Gamma_2, \Gamma_3, w : B \wp C \vdash \text{let-pat } w (y \wp -) \Delta_1 \mid \text{let-pat } w (- \wp z) \Delta_2 \\
\hline
\text{CUT} \\
\hline
\Gamma_1, \Gamma, \Gamma_2, \Gamma_3, w : B \wp C \vdash \Delta \mid [t/x](\text{let-pat } w (y \wp -) \Delta_1) \mid [t/x](\text{let-pat } w (- \wp z) \Delta_2)
\end{array}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2, y : B \vdash \Delta_1}}{\Gamma_1, \Gamma, \Gamma_2, y : B \vdash \Delta \mid [t/x]\Delta_1} \text{CUT} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, z : C \vdash \Delta_2}}{\Gamma_1, \Gamma, \Gamma_2, \Gamma_3, w : B \wp C \vdash \text{let-pat } w (y \wp -) \Delta \mid \text{let-pat } w (y \wp -) [t/x]\Delta_1 \mid \text{let-pat } w (- \wp z) \Delta_2} \text{PARL}$$

First, by inspection of the previous proofs we can see that  $y \notin \text{FV}(\Delta)$  and  $x \notin \text{FV}(\Delta_2)$ . Thus,  $\text{let-pat } w (y \wp -) \Delta = \Delta$ , and  $[t/x](\text{let-pat } w (- \wp z) \Delta_2) = \text{let-pat } w (- \wp z) \Delta_2$ . It suffices to show that  $[t/x](\text{let-pat } w (y \wp -) \Delta_1) = \text{let-pat } w (y \wp -) [t/x]\Delta_1$  but this follows by distributing the substitution into the let-pat, and then simplifying using the fact that  $w \neq x$ .

Case: secondary hypothesis: left introduction of par (second case). The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1, y : B \vdash \Delta_1} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_2, x : A, \Gamma_3, z : C \vdash \Delta_2}}{\Gamma_1, \Gamma_2, x : A, \Gamma_3, w : B \wp C \vdash \text{let-pat } w (y \wp -) \Delta_1 \mid \text{let-pat } w (- \wp z) \Delta_2} \text{PARL}}{\Gamma_1, \Gamma_2, \Gamma, \Gamma_3, w : B \wp C \vdash \Delta \mid [t/x](\text{let-pat } w (y \wp -) \Delta_1) \mid [t/x](\text{let-pat } w (- \wp z) \Delta_2)} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1, y : B \vdash \Delta_1} \quad \frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_2, x : A, \Gamma_3, z : C \vdash \Delta_2}}{\Gamma_2, \Gamma, \Gamma_3, z : C \vdash \Delta \mid [t/x]\Delta_2} \text{CUT}}{\Gamma_1, \Gamma_2, \Gamma, \Gamma_3, w : B \wp C \vdash \text{let-pat } w (y \wp -) \Delta_1 \mid \text{let-pat } w (- \wp z) \Delta \mid \text{let-pat } w (- \wp z) [t/x]\Delta_2} \text{PARL}$$

Similar to the previous case.

Case: secondary hypothesis: left introduction of implication (first case). The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2 \vdash t_1 : B \mid \Delta_1} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, y : C \vdash \Delta_2}}{\Gamma_1, x : A, \Gamma_2, \Gamma_3, z : B \multimap C \vdash \Delta_1 \mid [z t_1/y]\Delta_2} \text{IMPL}}{\Gamma_1, \Gamma, \Gamma_2, \Gamma_3, z : B \multimap C \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x][z t_1/y]\Delta_2} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2 \vdash t_1 : B \mid \Delta_1}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]t_1 : B \mid [t/x]\Delta_1} \text{CUT} \quad \frac{\frac{\pi_3}{\vdots}}{\Gamma_3, y : C \vdash \Delta_2}}{\Gamma_1, \Gamma, \Gamma_2, \Gamma_3, z : B \multimap C \vdash \Delta \mid [t/x]\Delta_1 \mid [z ([t/x]t_1)/y]\Delta_2} \text{IMPL}$$

By inspection of the above derivations we can see that  $x \notin \text{FV}(\Delta_2)$ , and hence, by this fact and substitution distribution (Lemma 1) we know  $[t/x][z t_1/y]\Delta_2 = [([t/x]z) ([t/x]t_1)/y][t/x]\Delta_2 = [z ([t/x]t_1)/y]\Delta_2$ .

Case: secondary hypothesis: left introduction of implication (second case). The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_1 \vdash t_1 : B \mid \Delta_1} \quad \frac{\pi_3}{\vdots}}{\Gamma_1, \Gamma_2, x : A, \Gamma_3, y : C \vdash \Delta_2} \text{IMPL}}{\Gamma_1, \Gamma_2, \Gamma, \Gamma_3, z : B \multimap C \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x][z t_1/y]\Delta_2} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1 \vdash t_1 : B \mid \Delta_1} \quad \frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\pi_3}{\vdots}}{\Gamma_2, x : A, \Gamma_3, y : C \vdash \Delta_2} \text{CUT}}{\frac{\Gamma_1, \Gamma_2, \Gamma, \Gamma_3, z : B \multimap C \vdash \Delta_1 \mid [z t_1/y]\Delta \mid [z t_1/y][t/x]\Delta_2}{\Gamma_1, \Gamma_2, \Gamma, \Gamma_3, z : B \multimap C \vdash [z t_1/y]\Delta \mid \Delta_1 \mid [z t_1/y][t/x]\Delta_2} \text{IMPL}}{\text{SERIES OF EXCHANGES}}$$

By inspection of the above proofs we can see that  $y \notin \text{FV}(\Delta)$ . Thus,  $[z t_1/y]\Delta = \Delta$ . The same can be said for the variable  $x$  and context  $\Delta_1$ , and hence,  $[t/x]\Delta_1 = \Delta_1$ . Finally, by inspection of the above proofs  $x \notin \text{FV}(t_1)$  and so by substitution distribution (Lemma 1) we know  $[t/x][z t_1/y]\Delta_2 = [z t_1/y][t/x]\Delta_2$ .

Case: secondary hypothesis: left introduction of implication (third case). The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_1 \vdash t_1 : B \mid \Delta_1} \quad \frac{\pi_3}{\vdots}}{\Gamma_1, \Gamma_2, z : B \multimap C, \Gamma_3, x : A \vdash \Delta_1 \mid [z t_1/y]\Delta_2} \text{IMPL}}{\Gamma_1, \Gamma_2, z : B \multimap C, \Gamma_3, \Gamma \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x][z t_1/y]\Delta_2} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_2}{\vdots}}{\Gamma_1 \vdash t_1 : B \mid \Delta_1} \quad \frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\pi_3}{\vdots}}{\Gamma_2, y : C, \Gamma_3, x : A \vdash \Delta_2} \text{CUT}}{\frac{\Gamma_1, \Gamma_2, z : B \multimap C, \Gamma_3, \Gamma \vdash \Delta \mid [t/x]\Delta_2}{\Gamma_1, \Gamma_2, z : B \multimap C, \Gamma_3, \Gamma \vdash \Delta \mid [z t_1/y]\Delta \mid [z t_1/y][t/x]\Delta_2} \text{IMPL}}{\Gamma_1, \Gamma_2, z : B \multimap C, \Gamma_3, \Gamma \vdash [z t_1/y]\Delta \mid \Delta_1 \mid [z t_1/y][t/x]\Delta_2} \text{SERIES OF EXCHANGES}$$

Similar to the previous case.

Case: secondary hypothesis: right introduction of implication. The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2, y : B \vdash t_1 : C \mid \Delta_1} \quad y \notin \text{FV}(\Delta_1)}{\Gamma_1, x : A, \Gamma_2 \vdash \lambda y. t_1 : B \multimap C \mid \Delta_1} \text{IMPR}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x](\lambda y. t_1) : B \multimap C \mid [t/x]\Delta_1} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2, y : B \vdash t_1 : C \mid \Delta_1} \text{CUT}}{\frac{\Gamma_1, \Gamma, \Gamma_2, y : B \vdash \Delta \mid [t/x]t_1 : C \mid [t/x]\Delta_1}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid \lambda y. [t/x]t_1 : B \multimap C \mid [t/x]\Delta_1} \text{IMPR}}$$



Clearly,  $[t/x](\lambda y.t_1) = \lambda y.[t/x]t_1$ .

Case: secondary hypothesis: left introduction of tensor unit. The proof

$$\frac{\frac{\pi_1}{\vdots} \quad \frac{\frac{\pi_2}{\vdots}}{G_1, x : A, G_2 \vdash \Delta_1} \text{IL}}{G \vdash t : A \mid \Delta \quad G_1, x : A, G_2, y : \top \vdash \text{let } y \text{ be } * \text{ in } \Delta_1} \text{CUT}}{G_1, G, G_2, y : \top \vdash \Delta \mid [t/x](\text{let } y \text{ be } * \text{ in } \Delta_1)} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{G \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{G_1, x : A, G_2 \vdash \Delta_1} \text{CUT}}{G_1, G, G_2 \vdash \Delta \mid [t/x]\Delta_1} \text{IL}}{G_1, G, G_2, y : \top \vdash \text{let } y \text{ be } * \text{ in } \Delta \mid \text{let } y \text{ be } * \text{ in } [t/x]\Delta_1} \text{IL}$$

It suffices to show that  $\Delta = \text{let } y \text{ be } * \text{ in } \Delta$  and  $[t/x](\text{let } y \text{ be } * \text{ in } \Delta_1) = \text{let } y \text{ be } * \text{ in } [t/x]\Delta_1$ . Without loss of generality suppose  $\Delta = t : B, \Delta'$ . We know that it must be the case that  $y \notin \text{FV}(t)$ , and we know that  $[y/z]t = t$  when  $z \notin \text{FV}(t)$ . Then by  $\text{EQ\_ETA2I}$  we have  $t = \text{let } y \text{ be } * \text{ in } t$ . This argument can be repeated for any other term in  $\Delta'$ . Thus,  $\Delta = \text{let } y \text{ be } * \text{ in } \Delta$ . It is easy to see that  $[t/x](\text{let } y \text{ be } * \text{ in } \Delta_1) = \text{let } y \text{ be } * \text{ in } [t/x]\Delta_1$  using the rule  $\text{EQ\_NATI}$ .

Case: secondary hypothesis: right introduction of par unit. The proof

$$\frac{\frac{\pi_1}{\vdots} \quad \frac{\frac{\pi_2}{\vdots}}{G_1, x : A, G_2 \vdash \Delta_1} \text{PR}}{G \vdash t : A \mid \Delta \quad G_1, x : A, G_2 \vdash \circ : \perp \mid \Delta_1} \text{CUT}}{G_1, G, G_2 \vdash \Delta \mid [t/x]\circ : \perp \mid [t/x]\Delta_1} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{G \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{G_1, x : A, G_2 \vdash \Delta_1} \text{CUT}}{G_1, G, G_2 \vdash \Delta \mid [t/x]\Delta_1} \text{PR}}{\frac{G_1, G, G_2 \vdash \circ : \perp \mid \Delta \mid [t/x]\Delta_1}{G_1, G, G_2 \vdash \Delta \mid \circ : \perp \mid [t/x]\Delta_1} \text{SERIES OF EXCHANGES}} \text{PR}$$

Clearly,  $[t/x]\circ = \circ$ .

Case: secondary hypothesis: left introduction of exchange. The proof

$$\frac{\frac{\pi_1}{\vdots} \quad \frac{\frac{\pi_2}{\vdots}}{G_1, x : A, G_2, w : B, y : C, G_3 \vdash \Delta_1} \text{EXL}}{G \vdash t : A \mid \Delta \quad G_1, x : A, G_2, y : C, w : B, G_3 \vdash \Delta_1} \text{CUT}}{G_1, G, G_2, y : C, w : B, G_3 \vdash \Delta \mid [t/x]\Delta_1} \text{CUT}$$

transforms into the proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2, w : B, y : C, \Gamma_3 \vdash \Delta_1}}{\Gamma_1, \Gamma, \Gamma_2, w : B, y : C, \Gamma_3 \vdash \Delta \mid [t/x]\Delta_1} \text{CUT}}{\Gamma_1, \Gamma, \Gamma_2, y : C, w : B, \Gamma_3 \vdash \Delta \mid [t/x]\Delta_1} \text{EXL}$$

Clearly, all terms are equivalent.

Case: secondary hypothesis: right introduction of exchange. The proof

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2 \vdash \Delta_1 \mid t_1 : B \mid t_2 : C \mid \Delta_2} \text{EXR}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x]t_2 : C \mid [t/x]t_1 : B \mid [t/x]\Delta_2} \text{CUT}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x]t_2 : C \mid [t/x]t_1 : B \mid [t/x]\Delta_2} \text{CUT}$$

is transformed into

$$\frac{\frac{\frac{\pi_1}{\vdots}}{\Gamma \vdash t : A \mid \Delta} \quad \frac{\frac{\pi_2}{\vdots}}{\Gamma_1, x : A, \Gamma_2 \vdash \Delta_1 \mid t_1 : B \mid t_2 : C \mid \Delta_2} \text{CUT}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x]t_1 : B \mid [t/x]t_2 : C \mid [t/x]\Delta_2} \text{EXR}}{\Gamma_1, \Gamma, \Gamma_2 \vdash \Delta \mid [t/x]\Delta_1 \mid [t/x]t_2 : C \mid [t/x]t_1 : B \mid [t/x]\Delta_2} \text{EXR}$$

Clearly, all terms are equivalent.

**Corollary 1 (Cut-Elimination).** *Cut-elimination holds for FILL.*

## 4 Full LNL Models

One of the difficult questions considering the categorical models of linear logic was how to model Girard's exponential,  $!$ , which is read "of course". The  $!$  modality can be used to translate intuitionistic logic into intuitionistic linear logic, and so the correct categorical interpretation of  $!$  should involve a relationship between a cartesian closed category, and the model of intuitionistic linear logic.

de Paiva gave some of the first categorical models of both classical and intuitionistic linear logic in her thesis [8]. She showed that a particular dialectica category called  $\text{Dial}_2(\text{Sets})$  is a model of FILL where  $!$  is interpreted as a comonad which produces natural comonoids, see page 76 of [8].

**Definition 4.** *The category  $\text{Dial}_2(\text{Sets})$  consists of*

- objects that are triples,  $A = (U, X, \alpha)$ , where  $U$  and  $X$  are sets, and  $\alpha \subseteq U \times X$  is a relation, and
- maps that are pairs  $(f, F) : (U, X, \alpha) \longrightarrow (V, Y, \beta)$  where  $f : U \longrightarrow V$  and  $F : Y \longrightarrow X$  such that
  - For any  $u \in U$  and  $y \in Y$ ,  $\alpha(u, F(y))$  implies  $\beta(f(u), y)$ .

Suppose  $A = (U, X, \alpha)$ ,  $B = (V, Y, \beta)$ , and  $C = (W, Z, \gamma)$ . Then identities are given by  $(\text{id}_U, \text{id}_X) : A \longrightarrow A$ . The composition of the maps  $(f, F) : A \longrightarrow B$  and  $(g, G) : B \longrightarrow C$  is defined as  $(f; g, G; F) : A \longrightarrow C$ .

In her thesis de Paiva defines a particular class of dialectica categories called  $GC$  over a base category  $C$ , see page 41 of [8]. The category  $\text{Dial}_2(\text{Sets})$  defined above can be seen as an instantiation of  $GC$  by setting  $C$  to be the category  $\text{Sets}$  of sets and functions between them. This model is a non-trivial model (all four units of the multiplicative and additive disjunction are different objects in the category), and does not model classical logic; see [8] page 58.

Seely gave a different, syntactic categorical model that confirmed that the of-course exponential should be modeled by a comonad [22]. However, Seely's model turned out to be unsound, as pointed out by Bierman [3]. This then prompted Bierman, Hyland, de Paiva, and Benton to define another categorical model called linear categories (Definition 5) that are sound, and also model  $!$  using a monoidal comonad [3].

**Definition 5.** A *linear category*,  $\mathcal{L}$ , consists of:

- A symmetric monoidal closed category  $\mathcal{L}$ ,
- A symmetric monoidal comonad  $(!, \epsilon, \delta, m_{A,B}, m_I)$  such that
  - For every free  $!$ -coalgebra  $(!A, \delta_A)$  there are two distinguished monoidal natural transformations  $e_A : !A \rightarrow I$  and  $d_A : !A \rightarrow !A \otimes !A$  which form a commutative comonoid and are coalgebra morphisms.
  - If  $f : (!A, \delta_A) \rightarrow (!B, \delta_B)$  is a coalgebra morphism between free coalgebras, then it is also a comonoid morphism.

This definition is the one given by Bierman in his thesis, see [3] for full definitions.

Intuitionistic logic can be interpreted in a linear category as a full subcategory of the category of  $!$ -coalgebras for the comonad, see proposition 17 of [3].

Benton gave a more balanced view of linear categories called LNL models.

**Definition 6.** A *linear/non-linear model (LNL model)* consists of

- a cartesian closed category  $(\mathcal{C}, 1, \times, \Rightarrow)$ ,
- a SMCC  $(\mathcal{L}, I, \otimes, \dashv)$ , and
- a pair of symmetric monoidal functors  $(G, n) : \mathcal{L} \rightarrow \mathcal{C}$  and  $(F, m) : \mathcal{C} \rightarrow \mathcal{L}$  between them that form a symmetric monoidal adjunction with  $F \dashv G$ .

See Benton, [2], for the definitions of symmetric monoidal functors and adjunctions.

A non-trivial consequence of the definition of a LNL model is that the  $!$  modality can indeed be interpreted as a monoidal comonad. Suppose  $(\mathcal{L}, \mathcal{C}, F, G)$  is a LNL model. Then the comonad is given by  $(!, \epsilon : ! \rightarrow \text{Id}, \delta : ! \rightarrow !!)$  where  $! = FG$ ,  $\epsilon$  is the counit of the adjunction and  $\delta$  is the natural transformation  $\delta_A = F(\eta_{G(A)})$ , see page 15 of [2]. We recall the following result from Benton [2]:

**Theorem 2 (LNL Models and Linear Categories).**

- i. (Section 2.2.1 of [2]) Every LNL model is a linear category.
- ii. (Section 2.2.2 of [2]) Every linear category is a LNL model.

*Proof.* The proof of part i. is a matter of checking that each part of the definition of a linear category can be constructed using the definition of a LNL model. See lemmata 3-7 of [2].

As for the proof of part ii. Given a linear category we have a SMCC and so the difficulty of proving this result is constructing the CCC and the adjunction between both parts of the model. Suppose  $\mathcal{L}$  is a linear category. Benton constructs the CCC out of the full subcategory of Eilenberg-Moore category  $\mathcal{L}^!$  whose objects are exponentiable coalgebras denoted  $\text{Exp}(\mathcal{L}^!)$ . He shows that this subcategory is cartesian closed, and contains the (co)Kleisli category,  $\mathcal{L}_!$ , Lemma 11 on page 23 of [2]. The required adjunction  $F : \text{Exp}(\mathcal{L}^!) \rightarrow \mathcal{L} : G$  can be defined using the adjunct functors  $F(A, h_A) = A$  and  $G(A) = (!A, \delta_A)$ , see lemmata 13 - 16 of [2].

Next we show that the category  $\text{Dia}_2(\text{Sets})$  is a full version of a linear category. First, we extend the definitions of linear categories and LNL models to be equipped with the necessary categorical structure to model par and its unit.

**Definition 7.** A *full linear category*,  $\mathcal{L}$ , consists of a linear category  $(\mathcal{L}, \top, \otimes, \dashv, !A, e_A, d_A)$ , a symmetric monoidal structure on  $L$ ,  $(\perp, \wp)$ , and distribution natural transformations  $\text{dist}_1 : A \otimes (B \wp C) \rightarrow (A \otimes B) \wp C$  and  $\text{dist}_2 : (A \wp B) \otimes C \rightarrow A \wp (B \otimes C)$ . The distributors must satisfy several coherence conditions which can all be found in [7].

**Definition 8.** A *full linear/non-linear model (full LNL model)* consists of a LNL model  $(\mathcal{L}, \mathcal{C}, F, G)$ , and a symmetric monoidal structure on  $L$ ,  $(\perp, \wp)$ , as above.

First we show that  $\text{Dial}_2(\mathbf{Sets})$  is a full linear category, and then using the proof by Benton that linear categories are LNL models we obtain that  $\text{Dial}_2(\mathbf{Sets})$  is a full LNL model. In order for this to work we need to know that  $\text{Dial}_2(\mathbf{Sets})$  has a symmetric *monoidal* comonad  $(!, \epsilon, \delta, m_{A,B}, m_I)$ . At the time of de Paiva's thesis it was not known that the comonad modeling the of-course exponential needed to be monoidal. We were able to show that the maps  $m_{A,B}$  and  $m_I$  exist in the more general setting of dialectica categories, and thus, these maps exist in  $\text{Dial}_2(\mathbf{Sets})$ . Intuitively, given two objects  $A = (X, U, \alpha)$  and  $B = (V, Y, \beta)$  of  $\text{Dial}_2(\mathbf{Sets})$  the map  $m_{A,B}$  is defined as the pair  $(\text{id}_{U \times V}, F)$ , where  $F = (F_1, F_2)$ ,  $F_1 : (U \times V) \Rightarrow (V \Rightarrow X)^* \longrightarrow V \Rightarrow (U \Rightarrow X^*)$  and  $F_2 : (U \times V) \Rightarrow (U \Rightarrow Y)^* \longrightarrow U \Rightarrow (V \Rightarrow Y^*)$ . The maps  $F_1$  and  $F_2$  build the sequence of all the results of applying each function in the input sequence to the input coordinate.

We can now show that the following holds.

**Lemma 3.** *The category  $\text{Dial}_2(\mathbf{Sets})$  is a full linear category.*

*Proof.* This proof holds by constructing each piece of a full linear category using the structure of  $\text{Dial}_2(\mathbf{Sets})$ <sup>4</sup>. We use some notation to make it easier to define and use functions over sequences. Given a function  $g : A \longrightarrow X^*$  we will denote taking the  $i$ th projection of the sequence returned by  $g(a)$  for some  $a \in A$  by  $g(a)_i$ . To construct set-theoretic anonymous functions we use  $\lambda$ -notation. Lastly, we often use let-expressions to pattern match on sequences. For example, let  $(x_1, \dots, x_i) = g(a)$  in  $(f(x_1), \dots, f(x_i))$ .

First, we must construct a linear category. It is well known that  $\mathbf{Sets}$  is a CCC, and in fact, locally cartesian closed, and so by using the results of de Paiva's thesis we can easily see that  $\text{Dial}_2(\mathbf{Sets})$  is symmetric monoidal closed:

- (Definition 7, page 43 of [8]). Suppose  $A, B \in \text{Obj}(\text{Dial}_2(\mathbf{Sets}))$ . Then there are sets  $X, Y, V$ , and  $U$ , and relations  $\alpha \subseteq U \times X$  and  $\beta \subseteq V \times Y$ , such that,  $A = (U, X, \alpha)$  and  $B = (V, Y, \beta)$ . The tensor product of  $A$  and  $B$  can now be defined by  $A \otimes B = (U \times V, (V \Rightarrow X) \times (U \Rightarrow Y), \alpha \otimes \beta)$ , where  $(- \Rightarrow -)$  is the internal hom of  $\mathbf{Sets}$ . We define  $((u, v), (f, g)) \in \alpha \otimes \beta$  if and only if  $(u, f(v)) \in \alpha$  and  $(v, g(u)) \in \beta$ . Suppose  $A = (U, X, \alpha)$ ,  $B = (V, Y, \beta)$ ,  $C = (W, Z, \gamma)$ , and  $D = (S, T, \delta)$  objects of  $\text{Dial}_2(\mathbf{Sets})$ , and  $m_1 = (f, F) : A \longrightarrow C$  and  $m_2 = (g, G) : B \longrightarrow D$  are maps of  $\text{Dial}_2(\mathbf{Sets})$ . Then the map  $m_1 \otimes m_2 : A \otimes B \longrightarrow C \otimes D$  is defined by  $(f \times g, F_\otimes)$  where  $f \times g$  is the ordinary cartesian product functor in  $\mathbf{Sets}$ , and we define  $F \otimes G$  as follows:

$$\begin{aligned} F_\otimes &: (S \Rightarrow Z) \times (W \Rightarrow T) \longrightarrow (V \Rightarrow X) \times (U \Rightarrow Y) \\ F_\otimes(h_1, h_2) &= (\lambda v. F(h_1(g(v))), \lambda u. G(h_2(f(u)))) \end{aligned}$$

It is straightforward to confirm the relation condition on maps for  $m_1 \otimes m_2$ .

- (Definition 7, page 44 of [8]). Suppose  $1 \in \text{Obj}(\mathbf{Sets})$  is the final object, and  $\text{id}_1 \subseteq 1 \times 1$ . Then we can define tensors unit by the object  $\top = (1, 1, \text{id}_1)$ .
- Suppose  $A = (U, X, \alpha)$  is an object of  $\text{Dial}_2(\mathbf{Sets})$ . Then the map  $\lambda_A : \top \otimes A \longrightarrow A$  is defined by  $(\hat{\lambda}_U, F_\lambda)$  where  $\hat{\lambda}_U$  is the left unitor for the cartesian product in  $\mathbf{Sets}$ ,  $F_\lambda(x) = (\diamond, \lambda y. x) : X \longrightarrow (U \Rightarrow 1) \times (1 \Rightarrow X)$ , and  $\diamond$  is the terminal arrow in  $\mathbf{Sets}$ . It is easy to see that both  $\hat{\lambda}_U$  and  $F_\lambda$  have inverses, and thus,  $\lambda_A$  has an inverse. It is straightforward to confirm the relation condition on maps for  $\lambda_A$  and its inverse.
- Suppose  $A = (U, X, \alpha)$  is an object of  $\text{Dial}_2(\mathbf{Sets})$ . Then the map  $\rho_A : A \otimes \top \longrightarrow A$  is defined similarly to  $\lambda_A$  given above.
- Suppose  $A = (U, X, \alpha) \in \text{Obj}(\text{Dial}_2(\mathbf{Sets}))$  and  $B = (V, Y, \beta) \in \text{Obj}(\text{Dial}_2(\mathbf{Sets}))$ . Then we define the map  $\beta_{A,B} : A \otimes B \longrightarrow B \otimes A$  by  $(\hat{\beta}_{U,V}, \hat{\beta}_{V \Rightarrow X, U \Rightarrow Y})$  where  $\hat{\beta}$  is the symmetry of the cartesian product in  $\mathbf{Sets}$ . Again, it is straightforward to see that  $\beta$  has an inverse, and the relation condition on maps is satisfied.
- Suppose  $A = (U, X, \alpha) \in \text{Obj}(\text{Dial}_2(\mathbf{Sets}))$ ,  $B = (V, Y, \beta) \in \text{Obj}(\text{Dial}_2(\mathbf{Sets}))$ , and  $C = (W, Z, \gamma) \in \text{Obj}(\text{Dial}_2(\mathbf{Sets}))$ . Then we define  $\alpha_{A,B,C} : (A \otimes B) \otimes C \longrightarrow A \otimes (B \otimes C)$  by  $(\hat{\alpha}_{U,V,W}, F_\alpha)$  where  $\hat{\alpha}_{U,V,W}$  is the associator for the cartesian product in  $\mathbf{Sets}$  and  $F_\alpha$  is defined as follows:

$$\begin{aligned} F_\alpha &: ((V \times W) \Rightarrow X) \times (U \Rightarrow ((W \Rightarrow Y) \times (V \Rightarrow Z))) \longrightarrow (W \Rightarrow ((V \Rightarrow X) \times (U \Rightarrow Y))) \times ((U \times V) \Rightarrow Z) \\ F_\alpha(h_1, h_2) &= (\lambda w. (\lambda v. h_1(v, w)), \lambda u. h_2(u)_1(w)), \lambda(u, v). h_2(u)_2(v) \end{aligned}$$

<sup>4</sup> This proof was formalized in the Agda proof assistant see the file <https://github.com/heades/cut-fill-agda/blob/master/FullLinCat.agda>

The inverse of  $\alpha_{A,B,C}$  is similar, and it is straightforward to confirm the relation condition on maps.

- (Definition 9, page 44 of [8]). Suppose  $A, B \in \text{Obj}(\text{Dial}_2(\text{Sets}))$ . Then there are sets  $X, Y, V$ , and  $U$ , and relations  $\alpha \subseteq U \times X$  and  $\beta \subseteq V \times Y$ , such that,  $A = (U, X, \alpha)$  and  $B = (V, Y, \beta)$ . Then we define the internal hom of  $\text{Dial}_2(\text{Sets})$  by  $A \multimap B = ((U \rightrightarrows V) \times (Y \rightrightarrows X), U \times Y, \alpha \rightrightarrows \beta)$ . We define  $((f, g), (u, y)) \in \alpha \rightrightarrows \beta$  if and only if whenever  $(u, g(y)) \in \alpha$ , then  $(f(u), y) \in \beta$ . The locally cartesian closed structure of  $\text{Sets}$  guarantees that for any two objects  $A, B \in \text{Dial}_2(\text{Sets})$  the internal hom  $A \multimap B \in \text{Dial}_2(\text{Sets})$  exists.

Using the constructions above  $\text{Dial}_2(\text{Sets})$  is a SMCC by Proposition 24 on page 46 of [8].

Next we define the symmetric monoidal comonad  $(!, \epsilon, \delta, m_{A,B}, m_I)$  of the linear category:

- (Section 4.5, on page 76 of [8]). The endofunctor  $! : \text{Dial}_2(\text{Sets}) \rightarrow \text{Dial}_2(\text{Sets})$  is defined as follows:
  - Objects. Suppose  $A = (U, X, \alpha) \in \text{Obj}(\text{Dial}_2(\text{Sets}))$ . Then we set  $!A = (U, U \rightrightarrows X^*, !\alpha)$ , where  $(u, f) \in !\alpha$  if and only if  $(u, f(u)_1) \in \alpha$  and  $\dots$  and  $(u, f(u)_i) \in \alpha$  where  $f(u)$  is a sequence of length  $i$ .
  - Morphisms. Suppose  $A = (U, X, \alpha) \in \text{Obj}(\text{Dial}_2(\text{Sets}))$ ,  $B = (V, Y, \beta) \in \text{Obj}(\text{Dial}_2(\text{Sets}))$ , and  $(f, F) : A \rightarrow B \in \text{Mor}(\text{Dial}_2(\text{Sets}))$ . Then we define  $!(f, F) = (f, !F) : !A \rightarrow !B$ , where  $!F(g) = \lambda x. F^*(g(f(x))) : V \rightrightarrows Y^* \rightarrow U \rightrightarrows X^*$ .
- (Section 4.5, page 77 of [8]). The endofunctor  $!$  defined above is the functor part of the comonad  $(!, \epsilon, \delta)$ . Suppose  $A = (U, X, \alpha) \in \text{Obj}(\text{Dial}_2(\text{Sets}))$ . Then the co-unit  $\epsilon : !A \rightarrow A$  is defined by  $\epsilon = (\text{id}_U, F_0)$  where  $F_0(x) = \lambda y. (x) : X \rightarrow U \rightrightarrows X^*$ . Furthermore, the co-multiplication  $\delta_A : !A \rightarrow !!A$  is defined by  $\delta_A = (\text{id}_U, F_1)$  where  $F_1(g) = \lambda u. (f_1(u) \circ \dots \circ f_i(u)) : U \rightrightarrows (U \rightrightarrows X^*)^* \rightarrow U \rightrightarrows X^*$  where  $g(u) = (f_1, \dots, f_i)$ .
- The following diagrams commute:

$$\begin{array}{ccc}
!A & \xrightarrow{\delta_A} & !!A \\
\delta_A \downarrow & & \downarrow !\delta_A \\
!!A & \xrightarrow{\delta_{!A}} & !!!A
\end{array}
\qquad
\begin{array}{ccc}
& !A & \\
& \swarrow & \searrow \\
!A & \xleftarrow{\epsilon_{!A}} & !!A \xrightarrow{! \epsilon_A} !A
\end{array}$$

We show the left most diagram commutes first. It suffices to show that  $\delta_a; !\delta_A = (\text{id}_U, !F_1; F_1) = (\text{id}_U, F_1; F_1)$ . Suppose  $g \in U \rightrightarrows (U \rightrightarrows X^*)^*$ . Then

$$\begin{aligned}
& F_1(F_1(g)) \\
&= F_1(\lambda u. g(u)_1(u) \circ \dots \circ g(u)_i(u)) \\
&= \lambda u. g(u)_1(u)_1(u) \circ \dots \circ g(u)_1(u)_j(u) \circ \dots \circ g(u)_i(u)_1(u) \circ \dots \circ g(u)_i(u)_k(u)
\end{aligned}$$

Consider the other direction in the diagram.

$$\begin{aligned}
F_1(!F_1(g)) &= F_1(\lambda x. F_1^*(g(x))) \\
&= \lambda u. F_1(g(u)_1)(u) \circ \dots \circ F_1(g(u)_i)(u)
\end{aligned}$$

Note that we have the following:

$$\begin{aligned}
F_1(g(u)_1)(u) &= g(u)_1(u)_1(u) \circ \dots \circ g(u)_1(u)_k(u) \\
&\vdots \\
F_1(g(u)_i)(u) &= g(u)_i(u)_1(u) \circ \dots \circ g(u)_i(u)_k(u)
\end{aligned}$$

Clearly, the above reasoning implies that  $F_1; F_1 = !F_1; F_1$ .

Now we prove that the second diagram commutes, but we break it into two. We define  $\delta_A; !\epsilon_A = (\text{id}_U, !F_0; F_1)$  where for any  $g \in U \rightrightarrows X^*$ ,

$$\begin{aligned}
(!F_0; F_1)(g) &= F_1(!F_0(g)) \\
&= F_1(\lambda u'. F_0^*(g(u'))) \\
&= F_1(\lambda u'. (\lambda y. (g(u')_1), \dots, (\lambda y. g(u')_i))) \\
&= \lambda u. (g(u)_1) \circ \dots \circ (g(u)_i) \\
&= g
\end{aligned}$$

and we can define  $\delta_A; \epsilon_{!A} = (\text{id}_U, F_0; F_1)$  where for any  $g \in U \Rightarrow X^*$ ,

$$\begin{aligned}
(F_0; F_1)(g) &= F_1(F_0(g)) \\
&= F_1(\lambda y. (g)) \\
&= \lambda u. g(u) \\
&= g
\end{aligned}$$

We can see by the reasoning above that  $!F_0; F_1 = F_0; F_1 = \text{id}_{U \Rightarrow X^*}$ .

- The monoidal natural transformation  $m_\top : \top \rightarrow !\top$  is defined by  $m_\top = (\text{id}_1, \lambda f. \star)$  where  $\star \in \top$ . It is easy to see that the relation condition on maps for  $\text{Dial}_2(\text{Sets})$  is satisfied. The following two diagrams commute:

$$\begin{array}{ccc}
\top & \xrightarrow{m_\top} & !\top \\
m_\top \downarrow & & \downarrow \delta_\top \\
!\top & \xrightarrow{!m_\top} & !!\top
\end{array}
\qquad
\begin{array}{ccc}
\top & & \\
m_\top \downarrow & \searrow & \\
!\top & \xrightarrow{\epsilon_\top} & \top
\end{array}$$

It is straightforward to see that the above two diagrams commute using the fact that the second coordinate of  $m_\top$  is a constant function.

- The monoidal natural transformation  $m_{A,B} : !A \otimes !B \rightarrow !(A \otimes B)$  is defined by  $m_{A,B} = (\text{id}_{U \times V}, F_2)$ . We need to define  $F_2$ , but two auxiliary functions are needed first:

$$\begin{aligned}
h_1 : (U \times V) \Rightarrow ((V \Rightarrow X) \times (U \Rightarrow Y))^* &\rightarrow (V \Rightarrow (U \Rightarrow X^*)) \\
h_1(g, v, u) &= (f_1(v), \dots, f_i(v)) \text{ where } g(u, v) = ((f_1, g_1), \dots, (f_i, g_i))
\end{aligned}$$

$$\begin{aligned}
h_2 : (U \times V) \Rightarrow ((V \Rightarrow X) \times (U \Rightarrow Y))^* &\rightarrow (U \Rightarrow (V \Rightarrow Y^*)) \\
h_2(g, u, v) &= (g_1(u), \dots, g_i(u)) \text{ where } g(u, v) = ((f_1, g_1), \dots, (f_i, g_i))
\end{aligned}$$

Then  $F_2(g) = (h_1(g), h_2(g))$ . In order for  $m_{A,B}$  to be considered a full fledged map in  $\text{Dial}_2(\text{Sets})$  we have to verify that the relation condition on maps is satisfied. Suppose  $(u, v) \in U \times V$  and  $g \in (U \times V) \Rightarrow ((V \Rightarrow X) \times (U \Rightarrow Y))^*$ , where  $g(u, v) = ((f_1, g_1), \dots, (f_i, g_i))$ . Then we know the following by definition:

$$\begin{aligned}
((u, v), F(g)) \in !\alpha \otimes !\beta &\text{ iff } ((u, v), (h_1(g), h_2(g))) \in !\alpha \otimes !\beta \\
&\text{ iff } (u, h_1(g)(v)) \in !\alpha \text{ and } (v, h_2(g)(u)) \in !\beta \\
&\text{ iff } (u, f_1(v)) \in \alpha \text{ and } \dots \text{ and } (u, f_i(v)) \text{ and} \\
&\quad (v, g_1(u)) \in \beta \text{ and } \dots \text{ and } (v, g_i(u))
\end{aligned}$$

and

$$\begin{aligned}
((u, v), g) \in !(\alpha \otimes \beta) &\text{ iff } ((u, v), (f_1, g_1)) \in \alpha \otimes \beta \text{ and } \dots \text{ and} \\
&\quad ((u, v), (f_i, g_i)) \in \alpha \otimes \beta \\
&\text{ iff } (u, f_1(v)) \in \alpha \text{ and } (v, g_1(u)) \in \beta \text{ and } \dots \text{ and} \\
&\quad (u, f_i(v)) \in \alpha \text{ and } (v, g_i(u)) \in \beta
\end{aligned}$$

The previous definitions imply that  $((u, v), F(g)) \in !\alpha \otimes !\beta$  implies  $((u, v), g) \in !(\alpha \otimes \beta)$ . Thus,  $m_{A,B}$  is a map in  $\text{Dial}_2(\text{Sets})$ .

At this point we show that the following diagrams commute:

$$\begin{array}{ccc}
!T \otimes !A & \xrightarrow{m_{T,A}} & !(T \otimes A) \\
\uparrow m_{T \otimes \text{id}_{!A}} & \text{A} & \downarrow !\lambda_A \\
T \otimes !A & \xrightarrow{\lambda_{!A}} & !A
\end{array}
\qquad
\begin{array}{ccc}
!A \otimes !T & \xrightarrow{m_{A,T}} & !(A \otimes T) \\
\uparrow \text{id}_{!A} \otimes m_T & \text{B} & \downarrow !\rho_A \\
!A \otimes T & \xrightarrow{\rho_{!A}} & !A
\end{array}$$

$$\begin{array}{ccc}
!A \otimes !B & \xrightarrow{m_{A,B}} & !(A \otimes B) \\
\downarrow \beta & \text{C} & \downarrow !\beta \\
!B \otimes !A & \xrightarrow{m_{B,A}} & !(B \otimes A)
\end{array}$$

$$\begin{array}{ccccc}
(!A \otimes !B) \otimes !C & \xrightarrow{m_{A,B} \otimes \text{id}_{!C}} & !(A \otimes B) \otimes !C & \xrightarrow{m_{A \otimes B, C}} & !((A \otimes B) \otimes C) \\
\downarrow \alpha_{!A, !B, !C} & & \text{D} & & \downarrow !\alpha_{A, B, C} \\
!A \otimes (!B \otimes !C) & \xrightarrow{\text{id}_{!A} \otimes m_{B,C}} & !A \otimes !(B \otimes C) & \xrightarrow{m_{A, B \otimes C}} & !(A \otimes (B \otimes C))
\end{array}$$

We first prove that diagram A commutes, and then diagrams B, and C will commute using similar reasoning. Following this we show that diagram D commutes. It suffices to show that

$$(m_{T \otimes \text{id}_{!A}}; m_{T,A}; !\lambda_A = (\hat{\lambda}_U, \lambda g. F_\otimes(F_2(F_\lambda(g)))) = (\hat{\lambda}_U, \lambda g. (\diamond, \lambda t. \lambda u. g(u))).$$

Suppose  $g \in U \Rightarrow X^*$ ,  $(\star, u) \in 1 \times U$ , and  $g(u) = (x_1, \dots, x_i)$ . Then

$$\begin{aligned}
F_\lambda(g)(\star, u) &= (\lambda x'. (\diamond, \lambda y. x'))^*(g(\hat{\lambda}_U(\star, u))) \\
&= (\lambda x'. (\diamond, \lambda y. x'))^*(g(u)) \\
&= (\lambda x'. (\diamond, \lambda y. x'))^*(x_1, \dots, x_i) \\
&= ((\diamond, \lambda y. x_1), \dots, (\diamond, \lambda y. x_i))
\end{aligned}$$

This implies that

$$F_\lambda(g) = \lambda(\star, u). \text{let } (x_1, \dots, x_i) = g(u) \text{ in } ((\diamond, \lambda y. x_1), \dots, (\diamond, \lambda y. x_i)).$$

Using this reasoning we can see the following:

$$\begin{aligned}
F_2(F_\lambda(g)) &= (\lambda u. \lambda t. \text{let } ((\diamond, \lambda y. x_1), \dots, (\diamond, \lambda y. x_i)) = F_\lambda(g)(t, u) \text{ in} \\
&\quad (\diamond(u), \dots, \diamond(u)), \\
&\quad \lambda t. \lambda u. \text{let } ((\diamond, \lambda y. x_1), \dots, (\diamond, \lambda y. x_i)) = F_\lambda(g)(t, u) \text{ in} \\
&\quad ((\lambda y. x_1)(t), \dots, (\lambda y. x_1)(t))) \\
&= (\lambda u. \lambda t. \text{let } ((\diamond, \lambda y. x_1), \dots, (\diamond, \lambda y. x_i)) = F_\lambda(g)(t, u) \text{ in} \\
&\quad (\star, \dots, \star), \\
&\quad \lambda t. \lambda u. \text{let } ((\diamond, \lambda y. x_1), \dots, (\diamond, \lambda y. x_i)) = F_\lambda(g)(t, u) \text{ in} \\
&\quad (x_1, \dots, x_1)) \\
&= (\lambda u. \lambda t. (\star, \dots, \star), \\
&\quad \lambda t. \lambda u. \text{let } ((\diamond, \lambda y. x_1), \dots, (\diamond, \lambda y. x_i)) = F_\lambda(g)(t, u) \text{ in} \\
&\quad (x_1, \dots, x_1)) \\
&= (\lambda u. \lambda t. (\star, \dots, \star), \lambda t. \lambda u. g(u))
\end{aligned}$$

Finally, the previous allows us to infer the following:

$$F_{\otimes}(F_2(F_{\lambda}(g))) = (\diamond, \lambda t. \lambda u. g(u))$$

Thus, we obtained our desired result.

We show that diagram D commutes by observing that

$$\begin{aligned} (m_{A,B} \otimes !\text{id}_C); m_{A \otimes B, C}; !\alpha_{A,B,C} &= (\text{id}, F_{\otimes}); (\text{id}, F_2); (\hat{\alpha}, !F_{\alpha}) \\ &= (\hat{\alpha}, !F_{\alpha}; F_2; F_{\otimes}) \\ &= (\hat{\alpha}, F_2; F_{\otimes}; F_{\alpha}) \\ &= (\hat{\alpha}, F_{\alpha}); (\text{id}, F_{\otimes}); (\text{id}, F_2) \end{aligned}$$

It suffices to show that  $!F_{\alpha}; F_2; F_{\otimes} = F_2; F_{\otimes}; F_{\alpha}$ :

$$\begin{aligned} (!F_{\alpha}; F_2; F_{\otimes})(g) &= F_{\otimes}(F_2(!F_{\alpha}(g))) \\ &= F_{\otimes}(F_2(\lambda x. F_{\alpha}^*(g(x)))) \end{aligned}$$

Suppose  $h_1 = \lambda v. \lambda u. \text{let}((f_1, g_1), \dots, (f_i, g_i)) = F_{\alpha}^*(g(u, v)) \text{ in } (f_1(v), \dots, f_i(v))$ ,

$h_2 = \lambda u. \lambda v. \text{let}((f_1, g_1), \dots, (f_i, g_i)) = F_{\alpha}^*(g(u, v)) \text{ in } (g_1(u), \dots, g_i(u))$ , and

$$\begin{aligned} F_{\alpha}^*(g(u, v)) &= \text{let}((f'_1, g'_1), \dots, (f'_j, g'_j)) = g(u, v) \text{ in} \\ &(\lambda w. (\lambda v'. f'_1(v', w), \lambda u. g'_1(u)_1(w)), \lambda(u, v'). g'_1(u)_2(v')), \dots, \\ &(\lambda w. (\lambda v'. f'_j(v', w), \lambda u. g'_j(u)_1(w)), \lambda(u, v'). g'_j(u)_2(v'))). \end{aligned}$$

Then we can simplify  $h_1$  and  $h_2$  as follows:

$$\begin{aligned} h_1 &= \lambda v. \lambda u. \text{let}((f'_1, g'_1), \dots, (f'_j, g'_j)) = g(u, v) \text{ in} \\ &((\lambda v'. f'_1(v', v), \lambda u'. g'_1(u')_1(v)), \dots, (\lambda v'. f'_j(v', v), \lambda u'. g'_j(u')_1(v))) \\ &\text{and} \\ h_2 &= \lambda u. \lambda v. \text{let}(u_1, u_2) = u \text{ in} \\ &\text{let}((f'_1, g'_1), \dots, (f'_j, g'_j)) = g((u_1, u_2), v) \text{ in} \\ &(g'_1(u_1)_2(u_2), \dots, g'_j(u_1)_2(u_2)) \end{aligned}$$

By the definition of  $F_2$  the previous reasoning implies:

$$\begin{aligned} F_{\otimes}(F_2(\lambda x. F_{\alpha}^*(g(x)))) &= F_{\otimes}(h_1, h_2) \\ &= (\lambda v. F_2(h_1(v)), h_2) \end{aligned}$$

Expanding the definition of  $F_2(h_1(v))$  in the above definitions yields:

$$F_2(h_1(v)) = (h'_1, h'_2)$$

where

$$\begin{aligned} h'_1 &= \lambda v''. \lambda u''. (f'_1(v'', v), \dots, f'_j(v'', v)) \\ h'_2 &= \lambda u''. \lambda v''. (g'_1(u')_1(v), \dots, g'_j(u')_1(v)) \end{aligned}$$

At this point we can see that

$$(\lambda v. F_2(h_1(v)), h_2) = (\lambda v. (h'_1, h'_2), h_2)$$

We now simplify  $F_2; F_{\otimes}; F_{\alpha}$ . We know by definition:

$$F_2(g) = (h''_1, h''_2)$$

where



$$\begin{aligned}
h_1'' &= \lambda v. \lambda u. \text{let } ((f_1', g_1'), \dots, (f_j', g_j')) = g(u, v) \text{ in} \\
&\quad \text{let } (v', v'') = v \text{ in } (f_1'(v', v''), \dots, f_k'(v', v'')) \\
\text{and} \\
h_2'' &= \lambda u. \lambda v. \text{let } ((f_1', g_1'), \dots, (f_j', g_j')) = g(u, v) \text{ in} \\
&\quad (g_1'(u), \dots, g_k'(u))
\end{aligned}$$

This implies that

$$\begin{aligned}
F_\alpha(F_\otimes(F_2(g))) &= F_\alpha(F_\otimes(h_1'', h_2'')) \\
&= F_\alpha(h_1'', \lambda u_4. F_2(h_2''(u_4))) \\
&= (\lambda w. (\lambda v. h_1''(v, w)), \lambda u. F_2(h_2''(u))_1(w)), \lambda(u, v). F_2(h_2''(u))_2(v)
\end{aligned}$$

Finally, by expanding the definition of  $F_2$  in the last line of the above reasoning we can see that

$$(\lambda v. (h_1', h_2'), h_2) = (\lambda w. (\lambda v. h_1''(v, w)), \lambda u. F_2(h_2''(u))_1(w)), \lambda(u, v). F_2(h_2''(u))_2(v))$$

modulo currying of set-theoretic functions.

- There are two coherence diagrams that  $m_{A,B}$  and  $\delta$  must adhere to. They are listed as follows:

$$\begin{array}{ccc}
!A \otimes !B & \xrightarrow{m_{A,B}} & !(A \otimes B) \\
\downarrow \epsilon_A \otimes \epsilon_B & \text{E} & \downarrow \epsilon_{A \otimes B} \\
A \otimes B & \xlongequal{\quad} & A \otimes B
\end{array}
\qquad
\begin{array}{ccccc}
!A \otimes !B & \xrightarrow{m_{A,B}} & & \xrightarrow{m_{A,B}} & !(A \otimes B) \\
\downarrow \delta_A \otimes \delta_B & & \text{F} & & \downarrow \delta_{A \otimes B} \\
!!A \otimes !!B & \xrightarrow{m_{!A,!B}} & !(!A \otimes !B) & \xrightarrow{!m_{A,B}} & !(A \otimes B)
\end{array}$$

Diagram E holds by simply expanding the definitions using an arbitrary input of a pair of functions. We now show diagram F commutes.

It suffices to show the following:

$$\begin{aligned}
m_{A,B}; \delta_{A \otimes B} &= (\text{id}_{U \times V}, F_1; F_2) \\
&= (\text{id}_{u \times V}, !F_2; F_2; F_\otimes) \\
&= (\delta_A \otimes \delta_B); m_{!A,!B}; !m_{A,B}
\end{aligned}$$

Suppose  $g \in (U \times V) \Rightarrow ((U \times V) \Rightarrow ((V \Rightarrow X) \times (U \Rightarrow Y))^*)^*$ . Then we know by the type of  $g$  and the definition of  $F_1$  it must be the case that  $F_1(g)$  first extracts all of the functions  $(f_1, \dots, f_i)$  returned by  $g(u, v)$  for arbitrary  $u \in U$  and  $v \in V$  – note that each  $f_i$  returns a sequence of pairs of functions,  $((f_i', g_i'), \dots, (f_j', g_j'))$  – then  $F_1(g)$  returns the concatenation of all of these sequences. Finally,  $F_2(F_1(g))$  returns two functions  $h_1(v, u)$  and  $h_2(u, v)$ , where  $h_1$  returns the sequence  $(f_i'(v), \dots, f_j'(v))$ , and  $h_2$  returns the sequence  $(g_i'(u), \dots, g_j'(u))$  from the sequence returned by  $F_1(g)$ . Note that each  $f_i'$  and  $g_i'$  returns a pair of functions.

Now consider applying  $!F_2; F_2; F_\otimes$  to  $g$ . The function  $!F_2$  will construct the function  $\lambda x. F_2^*(g(x))$  by definition, and  $F_2^*(g(x))$  will construct a sequence of pairs of functions  $((h_1', h_1''), \dots, (h_k', h_k''))$ . The function  $g$  as we saw above returns a sequence of functions,  $(f_1, \dots, f_i)$ , where each  $f_i$  returns a sequence of pairs of functions,  $((f_i', g_i'), \dots, (f_j', g_j'))$ . This tells us that by definition  $h_k'(v, u)$  will return the sequence  $(f_i'(v), \dots, f_j'(v))$  and  $h_k''(u, v)$  will construct the sequence  $(g_i'(u), \dots, g_j'(u))$ . Applying  $F_2$  to  $\lambda x. F_2^*(g(x))$  will construct two more functions  $t_1(v, u)$  and  $t_2(u, v)$  where the first returns the sequence of functions  $(h_1'(v), \dots, h_k'(v))$ , and the second returns  $(h_1''(u), \dots, h_k''(u))$ . Finally, applying the function  $F_\otimes$  to the pair  $(t_1, t_2)$  will result in a pair of functions

$$\begin{aligned}
(\lambda v. F_1(t_1(v)), \lambda u. F_1(t_2(u))) &= (\lambda v. \lambda u. h_1'(v)(u) \circ \dots \circ h_k'(v)(u), \\
&\quad \lambda u. \lambda v. h_1''(u)(v) \circ \dots \circ h_k''(u)(v)) \\
&= (\lambda v. \lambda u. (f_i'(v), \dots, f_j'(v)), \\
&\quad \lambda u. \lambda v. (g_i'(u), \dots, g_j'(u)))
\end{aligned}$$

We can now see that the pair  $(\lambda v.\lambda u.(f'_i(v), \dots, f'_j(v)), \lambda u.\lambda v.(g'_i(u), \dots, g'_k(u)))$  is indeed equivalent to the pair  $(h_1, h_2)$  given above, and thus, the diagram commutes.

Next we must show that whenever  $(!A, \delta)$  is a free comonoid, we have the distinguished natural transformations  $e_A : !A \rightarrow \top$  and  $d_A : !A \rightarrow !A \otimes !A$ . Suppose  $!A = (U, U \Rightarrow X^*)$  and  $(!A, \delta)$  is a free comonoid. Then we have the following definitions:

- (Proposition 53, page 77 of [8]). We define  $e_A : !A \rightarrow \top$  as the pair  $(\diamond, \lambda x.\lambda u.())$ , where  $\diamond$  is the terminal map on  $U$  and  $()$  is the empty sequence.
- (Proposition 53, page 77 of [8]). We define  $d_A : !A \rightarrow !A \otimes !A$  as the pair  $(\Delta, \theta)$  where  $\Delta : U \rightarrow U \times U$  is the diagonal map in **Sets**, and

$$\begin{aligned} \theta : ((U \times U) \Rightarrow X^*) \times ((U \times U) \Rightarrow X^*) &\rightarrow U \Rightarrow X^* \\ \theta(f, g) &= \lambda u.f(u, u) \circ g(u, u). \end{aligned}$$

The maps  $e_A$  and  $d_A$  must satisfy several coherence diagrams.

- We must show that the map  $e_A$  is a monoidal natural transformation. This requires that the following diagrams hold (for any arbitrary map  $f$ ):

$$\begin{array}{ccc} \begin{array}{ccc} !A & \xrightarrow{e_A} & \top \\ \downarrow f & \text{G} & \parallel \\ !B & \xrightarrow{e_B} & \top \end{array} & \begin{array}{ccc} \top & & \\ \downarrow m_I & \text{H} & \searrow \\ !\top & \xrightarrow{\epsilon_\top} & \top \end{array} & \begin{array}{ccc} !A \otimes !B & \xrightarrow{e_A \otimes e_B} & \top \otimes \top \\ \downarrow m_{A,B} & \text{I} & \downarrow \lambda \\ !(A \otimes B) & \xrightarrow{e_{A \otimes B}} & \top \end{array} \end{array}$$

Diagrams G and H follow easily by the definition of  $e_A$  and  $m_I$ . We now show that diagram I commutes. It suffices to show the following:

$$\begin{aligned} (e_A \otimes e_B); \lambda &= (\diamond_U \times \diamond_V, F_\otimes); (\hat{\lambda}_\top, F_\lambda) \\ &= ((\diamond_U \times \diamond_V); \hat{\lambda}_\top, F_\lambda; F_\otimes) \\ &= (\diamond_{U \times V}, F_\lambda; F_\otimes) \\ &= (\diamond_{U \times V}, F_2(\lambda u.())) \\ &= (\diamond_{U \times V}, (\lambda x.\lambda u.()); F_2) \\ &= (\text{id}_{U \times V}; \diamond_{U \times V}, (\lambda x.\lambda u.()); F_2) \\ &= (\text{id}_{U \times V}, F_2); (\diamond_{U \times V}, \lambda x.\lambda u.()) \\ &= m_{A,B}; e_{A \otimes B} \end{aligned}$$

It suffices to show  $F_\lambda; F_\otimes = F_2(\lambda u.())$ , but this easily follows by definition.

- The map  $d_A$  must be a monoidal natural transformation. This requires the following diagrams to commute:

$$\begin{array}{ccc}
!A & \xrightarrow{d_A} & !A \otimes !A \\
!f \downarrow & & \downarrow !f \otimes !f \\
!B & \xrightarrow{d_B} & !B \otimes !B
\end{array}
\quad \text{J}
\qquad
\begin{array}{ccc}
\top & \xrightarrow{\lambda^{-1}} & \top \otimes \top \\
m_{\top} \downarrow & & \downarrow m_{\top} \otimes m_{\top} \\
!\top & \xrightarrow{d_{\top}} & !\top \otimes !\top
\end{array}
\quad \text{K}$$
  

$$\begin{array}{ccc}
!A \otimes !B & \xrightarrow{d_A \otimes d_B} & (!A \otimes !A) \otimes (!B \otimes !B) & \xrightarrow{iso} & (!A \otimes !B) \otimes (!A \otimes !B) \\
m_{A,B} \downarrow & & \downarrow m_{A,B} \otimes m_{A,B} \\
!(A \otimes B) & \xrightarrow{d_{A \otimes B}} & !(A \otimes B) \otimes !(A \otimes B)
\end{array}
\quad \text{L}$$

Diagrams J and K follow easily from unfolding their definitions. We show that diagram L next. The morphism  $iso$  in  $\text{Dial}_2(\text{Sets})$  is an isomorphism that can be built out of the SMCC structure. For its definition in terms of the SMCC maps see footnote 9 on page 141 of [3], but we give a direct definition instead. We will need the following definitions:

$$\begin{aligned}
\hat{iso}((u, u'), (v, v')) &= ((u, v), (u', v')) \\
F_{iso}(f, g) &= (\lambda(v', v'').(\lambda u'.\lambda u''.f(u', v')_1(v'', u''), \lambda u'.\lambda u''.g(u', v')_1(v'', u'')), \\
&\quad \lambda(u', u'').(\lambda v'.\lambda v''.f(u', v')_2(u'', v''), \lambda v'.\lambda v''.g(u', v')_1(u'', v''))) \\
F_{iso}^{-1}(h_1, h_2) &= (\lambda(u, v).(\lambda v'.\lambda u'.h_1(v, v')_1(u, u'), \lambda u'.\lambda v'.h_2(u, u')_2(v, v')), \\
&\quad \lambda(u, v).(\lambda v'.\lambda u'.h_1(v, v')_2(u, u'), \lambda u'.\lambda v'.h_2(u, u')_2(v, v')))
\end{aligned}$$

We omit the proof that  $F_{iso}$  is an isomorphism, but it is straightforward. Now  $iso = (\hat{iso}, F_{iso})$ . It suffices to show the following:

$$\begin{aligned}
(d_A \otimes d_B); iso; (m_{A,B} \otimes m_{A,B}) &= (\Delta_U \times \Delta_V, F_{\otimes}); (\hat{iso}, F_{iso}); (\text{id}_{(U \times V) \times (U \times V)}, F_{\otimes}) \\
&= ((\Delta_U \times \Delta_V); \hat{iso}; \text{id}_{(U \times V) \times (U \times V)}, F_{\otimes}; F_{iso}; F_{\otimes}) \\
&= ((\Delta_U \times \Delta_V); \hat{iso}, F_{\otimes}; F_{iso}; F_{\otimes}) \\
&= (\Delta_{U \times V}, \Theta; F_2) \\
&= (\text{id}_{U \times V}; \Delta_{U \times V}, \Theta; F_2) \\
&= (\text{id}_{U \times V}, F_2); (\Delta_{U \times V}, \Theta) \\
&= m_{A,B}; d_{A,B}
\end{aligned}$$

At this point it suffices to show that  $F_{\otimes}; F_{iso}; F_{\otimes} = \Theta; F_2$ , but this follows using similar reasoning as above, because  $F_{iso}; F_{\otimes}$  will reorganize the streams obtained by applying  $g_1$  and  $g_2$ , and then the final  $F_{\otimes}$  combines these sequences using  $\Theta$ . However,  $F_2$  does the same reorganization, and then the streams are combined using  $\Theta$ .

- Suppose  $A \in \text{Obj}(\text{Dial}_2(\text{Sets}))$ . Then we must show that  $(!A, d_A, e_A)$  is a commutative comonoid, but this follows from the following diagrams:

$$\begin{array}{ccc}
& !A & \\
\rho^{-1} \swarrow & \downarrow d_A & \searrow \lambda^{-1} \\
!A \otimes T & !A \otimes !A & T \otimes !A \\
\text{id}_{!A} \otimes e_A \longleftarrow & & \longrightarrow e_A \otimes \text{id}_{!A} \\
\text{M} & & \text{N}
\end{array}
\qquad
\begin{array}{ccc}
!A & \xrightarrow{d_A} & !A \otimes !A \\
\parallel & & \downarrow \beta_{!A, !A} \\
!A & \xrightarrow{d_A} & !A \otimes !A \\
\text{O} & & 
\end{array}$$

$$\begin{array}{ccc}
!A & \xrightarrow{d_A} & !A \otimes !A \\
\downarrow d_A & & \downarrow \text{id}_{!A} \otimes d_A \\
!A \otimes !A & \xrightarrow{d_A \otimes \text{id}_{!A}} & (!A \otimes !A) \otimes !A \xrightarrow{\alpha} !A \otimes (!A \otimes !A) \\
\text{P} & & 
\end{array}$$

We prove that diagram N commutes, and then diagrams M and O will commute by similar reasoning. Following this we prove that diagram P commutes.

It suffices to show the following:

$$\begin{aligned}
d_A; (e_A \otimes \text{id}_{!A}) &= (\Delta, \Theta); (\diamond \times \text{id}_U, F_\otimes) \\
&= (\Delta; (\diamond \times \text{id}_U), F_\otimes; \Theta) \\
&= (\lambda u. (\diamond(u), u), F_\otimes; \Theta) \\
&= (\hat{\lambda}_U^{-1}, F_\otimes; \Theta) \\
&= (\hat{\lambda}^{-1}, F_{\lambda^{-1}}) \\
&= \lambda^{-1}
\end{aligned}$$

We can easily see that the following holds by definition:

$$\begin{aligned}
(F_\otimes; \Theta)(g_1, g_2) &= \Theta(F_\otimes(g_1, g_2)) \\
&= \Theta(\lambda v. \lambda u. (), \lambda u. g_2(\diamond(u))) \\
&= \lambda u. () \circ g_2(\diamond(u, u)) \\
&= \lambda u. g_2(\diamond(u, u)) \\
&= \lambda^{-1}(g_1, g_2)
\end{aligned}$$

Now we show that diagram P commutes. However, it is straightforward to show that the following holds:

$$\begin{aligned}
d_A; (\text{id}_{!A} \otimes d_A) &= (\Delta, \Theta); (\text{id}_U \times \Delta, F_\otimes) \\
&= (\Delta; (\text{id}_U \times \Delta), F_\otimes; \Theta) \\
&= (\Delta; (\text{id}_U \times \Delta), F_\alpha; F_\otimes; \Theta) \\
&= (\Delta; (\Delta \times \text{id}_U); \hat{\alpha}, F_\alpha; F_\otimes; \Theta) \\
&= (\Delta, \Theta); (\Delta \times \text{id}_U, F_\otimes); (\hat{\alpha}, F_\alpha) \\
&= d_A; (d_A \otimes \text{id}_{!A}); \alpha
\end{aligned}$$

We can see that  $F_\otimes; \Theta = F_\alpha; F_\otimes; \Theta$ , because the right-hand side does the same as the left-hand side, but first reorganizes and does it on the opposite association.

- The map  $e_A$  must be a coalgebra morphism which amounts to requiring that the following diagram commute:

$$\begin{array}{ccc}
!A & \xrightarrow{e_A} & T \\
\delta_A \downarrow & & \downarrow m_T \\
!!A & \xrightarrow{!e_A} & !T \\
\text{Q} & & 
\end{array}$$

Diagram Q commutes by unfolding definitions, and the fact that the second coordinate of  $e_A$  is a constant function.

- The map  $d_A$  must also be a coalgebra morphism, and hence, the following diagram must commute:

$$\begin{array}{ccccc}
 !A & \xrightarrow{\delta_A} & & \gg & !!A \\
 \downarrow d_A & & & & \downarrow !d_A \\
 !A \otimes !A & \xrightarrow{\delta_A \otimes \delta_A} & & \gg & !!A \otimes !!A \xrightarrow{m_{!A, !A}} & \gg & !(A \otimes !A)
 \end{array}$$

R

It suffices to show that the following holds:

$$\begin{aligned}
 \delta_A; !d_A &= (\text{id}_U, F_1); (\Delta, !\Theta) \\
 &= (\Delta, !\Theta; F_1) \\
 &= (\Delta; \text{id}_{U \times U}; \text{id}_{U \times U}, !\Theta; F_1) \\
 &= (\Delta; (\text{id}_U \times \text{id}_U); \text{id}_{U \times U}, !\Theta; F_1) \\
 &= (\Delta; (\text{id}_U \times \text{id}_U); \text{id}_{U \times U}, F_2; F_\otimes; \Theta) \\
 &= (\Delta, \Theta); (\text{id}_U \times \text{id}_U, F_\otimes); (\text{id}_{U \times U}, F_2) \\
 &= d_A; (\delta_A \otimes \delta_A); m_{!A, !A}
 \end{aligned}$$

Now consider the following:

$$\begin{aligned}
 !(\Theta; F_1)(g) &= F_1(!\Theta(g)) \\
 &= F_1(\lambda x. \Theta^*(g(x))) \\
 &= \lambda p. \text{let } (f_1, \dots, f_i) = \Theta^*(g(p)) \text{ in } f_1(p) \circ \dots \circ f_i(p)
 \end{aligned}$$

Furthermore, consider the following:

$$\begin{aligned}
 (F_2; F_\otimes; \Theta)(g) &= \Theta(F_\otimes(h_1, h_2)) \\
 &= \Theta(\lambda x. F_1(h_1(x)), \lambda y. F_1(h_2(y))) \\
 &= \lambda u. F_1(h_1(u, u)) \circ F_1(h_2(u, u))
 \end{aligned}$$

where

$$\begin{aligned}
 h_1(u, u') &= \text{let } (f_1, \dots, f_i) = g(u, u') \text{ in } (f_1(u), \dots, f_i(u)) \\
 h_2(u, u') &= \text{let } (f_1, \dots, f_i) = g(u, u') \text{ in } (f_1(u'), \dots, f_i(u'))
 \end{aligned}$$

At this point we can see that  $\Theta; F_1 = F_2; F_\otimes; \Theta$  by the previous reasoning and the definition of  $F_1$ .

- Finally, we show that when given a coalgebra morphism,  $f$ , between free coalgebras  $(!A, \delta_A)$  and  $(!B, \delta_B)$ , i.e. making the following diagram commute:

$$\begin{array}{ccc}
 !A & \xrightarrow{f} & !B \\
 \delta_A \downarrow & & \downarrow \delta_B \\
 !!A & \xrightarrow{!f} & !!B
 \end{array}$$

S

Then it must be the case that the following commutes:

$$\begin{array}{ccccc}
 \top & \xleftarrow{e_A} & !A & \xrightarrow{d_A} & !A \otimes !A \\
 \parallel & & \downarrow f & & \downarrow f \otimes f \\
 \top & \xleftarrow{e_B} & !B & \xrightarrow{d_B} & !B \otimes !B
 \end{array}$$

It is straightforward to show that the previous diagram commutes using the definitions of the respective morphisms and the assumption that  $f$  is a coalgebras morphism.

**Corollary 2.** *The category  $\text{Dial}_2(\text{Sets})$  is a full LNL model.*

*Proof.* This follows directly from the previous lemma and Theorem 2 which shows that linear categories are LNL models.

The point of these calculations is to show that the several different axiomatizations available for models for linear logic are consistent and that a model proved sound and complete according to Seely's definition (using the Seely isomorphisms  $!(A \times B) \cong !A \otimes !B$  and  $!1 \cong \top$  but adding to it monoidicity of the comonad) is indeed sound and complete as a LNL model too.

## 5 Tensorial Logic

Melliès and Tabareau introduced tensorial logic as a means of generalizing linear logic to a theory of tensor and a non-involutive negation called tensorial negation. That is, instead of an isomorphism  $A = \neg\neg A$  we have only a natural transformation  $A \rightarrow \neg\neg A$  [17]. Tensorial logic makes the claim that tensor and tensorial negation are more fundamental than tensor and negation defined via implication. This is at odds with FILL where implication is considered to be fundamental. In this section we show that multiplicative tensorial logic can be modeled by  $\text{Dial}_2(\text{Sets})$  (Lemma 5) by showing that tensorial negation arises as a simple property of the implication in any SMCC (Lemma 4). While this is expected (after all negation being defined in terms of implication into absurdity is one of the staples of intuitionism) we think it bolsters our claim that linear implication is a fundamental connective that should not be redefined in terms of the multiplicative disjunction par. In any case, any model of FILL can be seen as a model of multiplicative tensorial logic.

A categorical model of tensorial logic is a symmetric monoidal category with a tensorial negation.

**Definition 9.** *A **tensorial negation** on a symmetric monoidal category  $(\mathcal{C}, \otimes, I)$  is defined as a functor  $\neg : \mathcal{C} \rightarrow \mathcal{C}^{\text{op}}$  together with a family of bijections  $\phi_{A,B,C} : \text{Hom}_{\mathcal{C}}(A \otimes B, \neg C) \cong \text{Hom}_{\mathcal{C}}(A, \neg(B \otimes C))$  natural in  $A, B,$  and  $C$ . Furthermore, the following diagram must commute:*

$$\begin{array}{ccc}
\text{Hom}(A \otimes (B \otimes C), \neg D) & \xrightarrow{\text{Hom}(\alpha_{A,B,C}, \text{id}_{\neg D})} & \text{Hom}((A \otimes B) \otimes C, \neg D) \\
\downarrow \phi_{A,B \otimes C, D} & & \downarrow \phi_{A \otimes B, C, D} \\
& & \text{Hom}(A \otimes B, \neg(C \otimes D)) \\
& & \downarrow \phi_{A,B,C \otimes D} \\
\text{Hom}(A, \neg((B \otimes C) \otimes D)) & \xrightarrow{\text{Hom}(\text{id}_A, \neg\alpha_{B,C,D})} & \text{Hom}(A, \neg(B \otimes (C \otimes D)))
\end{array}$$

The most basic form of tensorial logic is called multiplicative tensorial logic and only consists of tensor and a tensorial negation. The model of multiplicative tensorial logic is called a dialogue category.

**Definition 10.** *A **dialogue category** is a symmetric monoidal category equipped with a tensorial negation.*

At this point we show that tensorial negation arises as a simple property of implication, as is traditional.

**Lemma 4.** *In any monoidal closed category,  $\mathcal{C}$ , there is a natural bijection  $\phi_{A,B,C,D} : \text{Hom}_{\mathcal{C}}(A \otimes B, C \multimap D) \cong \text{Hom}_{\mathcal{C}}(A, (B \otimes C) \multimap D)$ . Furthermore, the following diagram commutes:*

$$\begin{array}{ccc}
\text{Hom}(A \otimes (B \otimes C), D \multimap E) & \xrightarrow{\text{Hom}(\alpha_{A,B,C}, \text{id}_{D \multimap E})} & \text{Hom}((A \otimes B) \otimes C, D \multimap E) \\
\downarrow \phi_{A,B \otimes C, D, E} & & \downarrow \phi_{A \otimes B, C, D, E} \\
& & \text{Hom}(A \otimes B, (C \otimes D) \multimap E) \\
& & \downarrow \phi_{A,B,C \otimes D, E} \\
\text{Hom}(A, ((B \otimes C) \otimes D) \multimap E) & \xrightarrow{\text{Hom}(\text{id}_A, \alpha_{B,C,D \multimap E})} & \text{Hom}(A, (B \otimes (C \otimes D)) \multimap E)
\end{array}$$

*Proof.* Suppose  $\mathcal{C}$  is a monoidal closed category. Then we can define  $\phi(f : A \otimes B \multimap C \multimap D) = \text{cur}(\alpha^{-1}; \text{cur}^{-1}(f))$  and  $\phi^{-1}(g : A \multimap (B \otimes C) \multimap D) = \text{cur}(\alpha; \text{cur}^{-1}(g))$ . Clearly, these are mutual inverses, and hence,  $\phi$  is a bijection. Naturality of  $\phi$  easily follows.

Suppose  $f : A \otimes (B \otimes C) \multimap D \multimap E$ . The required diagram commutes by the following equational reasoning:

$$\begin{aligned}
\phi(\phi(\alpha; f)) &= \phi(\text{cur}(\alpha^{-1}; \text{cur}^{-1}(\alpha; f))) && \text{(Definition)} \\
&= \text{cur}(\alpha^{-1}; \text{cur}^{-1}(\text{cur}(\alpha^{-1}; \text{cur}^{-1}(\alpha; f)))) && \text{(Definition)} \\
&= \text{cur}(\alpha^{-1}; (\alpha^{-1}; \text{cur}^{-1}(\alpha; f))) && \text{(Inverses)} \\
&= \text{cur}((\alpha^{-1}; \alpha^{-1}); \text{cur}^{-1}(\alpha; f)) && \text{(Associativity)} \\
&= \text{cur}((\alpha^{-1}; \alpha^{-1}); (\alpha \otimes \text{id}); \text{cur}^{-1}(f)) && \text{(Naturality of cur)} \\
&= \text{cur}((\text{id} \otimes \alpha^{-1}); \alpha^{-1}; (\alpha^{-1} \otimes \text{id}); (\alpha \otimes \text{id}); \text{cur}^{-1}(f)) && \text{(Monoidal Pentagon)} \\
&= \text{cur}((\text{id} \otimes \alpha^{-1}); \alpha^{-1}; (\alpha^{-1}; \alpha \otimes \text{id}); \text{cur}^{-1}(f)) && \text{(Functorality)} \\
&= \text{cur}((\text{id} \otimes \alpha^{-1}); \alpha^{-1}; (\text{id} \otimes \text{id}); \text{cur}^{-1}(f)) && \text{(Inverses)} \\
&= \text{cur}((\text{id} \otimes \alpha^{-1}); \alpha^{-1}; \text{id}; \text{cur}^{-1}(f)) && \text{(Functorality)} \\
&= \text{cur}((\text{id} \otimes \alpha^{-1}); \alpha^{-1}; \text{cur}^{-1}(f)) && \text{(Identity)} \\
&= \text{cur}(\alpha^{-1}; \text{cur}^{-1}(f)); (\alpha^{-1} \multimap \text{id}) && \text{(Naturality of cur)} \\
&= \phi(f); (\alpha^{-1} \multimap \text{id}) && \text{(Definition)}
\end{aligned}$$

Any model of FILL contains the unit of  $\text{par}$ ,  $\perp$ , and thus, can be used to define the negation function  $\neg A := A \multimap \perp$ . Now replacing  $D$  and  $E$  in the previous result with  $\perp$  yields the definition of tensorial negation.

**Lemma 5.** *Dial<sub>2</sub>(Sets) is a model of multiplicative tensorial logic.*

*Proof.* We have already shown Dial<sub>2</sub>(Sets) to be a model of FILL, and thus, has a SMCC structure as well as the negation functor, and thus, by Lemma 4 has a tensorial negation<sup>5</sup>.

Extending a model of multiplicative tensorial logic with an exponential resource modality yields a model of full tensorial logic.

**Definition 11.** A **resource modality** on a symmetric monoidal category  $(\mathcal{C}, \otimes, I)$  is an adjunction with a symmetric monoidal category  $(\mathcal{M}, \otimes', I')$ :

$$\begin{array}{ccc}
& & \mathcal{C} \\
& \xrightarrow{F} & \\
\mathcal{M} & & \\
& \xleftarrow{G} & \\
& & \mathcal{C}
\end{array}$$

A resource modality is called an **exponential resource modality** if  $\mathcal{M}$  is cartesian where  $\otimes'$  is the product and  $I'$  is the terminal object.

A model of full tensorial logic is defined to be a model of multiplicative tensorial logic with an exponential resource modality. We now know that Dial<sub>2</sub>(Sets) is a model of multiplicative tensorial logic. By constructing the co-Kleisli category which consists of the !-coalgebras as objects, and happens to be cartesian, we can show that Dial<sub>2</sub>(Sets) is a model of full tensorial logic. The adjunction with the co-Kleisli category naturally arises from the proof that Dial<sub>2</sub>(Sets) is a full LNL model (Corollary 2).

**Lemma 6.** *The category Dial<sub>2</sub>(Sets) is a model of full tensorial logic.*

*Proof.* It suffices to show that there is an adjunction between Dial<sub>2</sub>(Sets) and a cartesian category. Define the category Dial<sub>2</sub>(Sets)<sub>!</sub> as follows:

- Take as objects  $(U, (U \Rightarrow X^*), \alpha_!)$  where  $U$  and  $X$  are sets, and  $\alpha \subseteq U \times (U \Rightarrow X^*)$ .

<sup>5</sup> We give a full proof in the formalization see the file <https://github.com/heades/cut-fill-agda/blob/master/Tensorial.agda>.

- Take as morphisms  $(f, F) : (U, (U \Rightarrow X^*), \alpha_1) \longrightarrow (V, (V \Rightarrow Y^*), \beta_1)$  where  $f : U \longrightarrow V$  and  $F : (V \Rightarrow Y^*) \longrightarrow (U \Rightarrow X^*)$  subject to the same condition on morphisms as  $\text{Dial}_2(\text{Sets})$ . Composition and identities are defined similarly to  $\text{Dial}_2(\text{Sets})$ .

Now we show that  $\text{Dial}_2(\text{Sets})_!$  is cartesian. Notice that  $\text{Dial}_2(\text{Sets})_!$  is a subcategory of  $\text{Dial}_2(\text{Sets})$ , and there is a functor  $J : \text{Dial}_2(\text{Sets}) \longrightarrow \text{Dial}_2(\text{Sets})_!$  which is defined equivalently to the endofunctor  $!$  from the proof of Lemma 3. In fact,  $\text{Dial}_2(\text{Sets})_!$  is the co-Kleisli category with objects free  $!$ -coalgebras and is cartesian closed [9]. However, we only need the fact that it is cartesian.

- Suppose  $A = (U, X, \alpha)$  and  $B = (V, Y, \beta)$  are objects of  $\text{Dial}_2(\text{Sets})$ . Then we define  $A \times B = (U \times V, (X + Y), \alpha \times \beta)$ , where  $((u, v), i) \in \alpha \times \beta$  iff when  $i \in X$ , then  $(u, i) \in \alpha$ , otherwise when  $i \in Y$ , then  $(v, i) \in \beta$ . Now the cartesian product in  $\text{Dial}_2(\text{Sets})_!$  is defined as  $J(A \times B)$ .
- The terminal object in  $\text{Dial}_2(\text{Sets})$  is defined by  $(\top, \perp, \alpha_\top)$  where  $\top$  and  $\perp$  are the terminal and initial objects in  $\text{Sets}$  respectively, and  $(x, y) \in \alpha_\top$  iff true. The proof that this is terminal, and is the unit to the cartesian product can be found in the formal development.
- Suppose  $A = (U, X, \alpha)$ ,  $B = (V, Y, \beta)$ , and  $C = (W, Z, \gamma)$  are objects of  $\text{Dial}_2(\text{Sets})$ . Then we define the following morphisms – the proofs of the morphism conditions have been omitted, but the details can be found in the formal development:
  - Define the first projection by  $(\pi_1, F_{\pi_1}) : J(A \times B) \longrightarrow J(A)$ , where  $\pi_1$  is the first projection in  $\text{Sets}$ , and  $F_{\pi_1}$  takes a function  $f : U \longrightarrow X^*$  and a pair  $(u, v) \in U \times V$ , and returns the sequence  $(X + Y)^*$  by mapping the first coproduct injection over  $f(u, v)$ .
  - The second projection is defined similar to the first,  $(\pi_2, F_{\pi_2}) : J(A \times B) \longrightarrow J(B)$ , but  $F_{\pi_2}$  maps the second projection instead of the first.
  - Suppose  $j_1 = (f, F) : J(C) \longrightarrow J(A)$  and  $j_2 = (g, G) : J(C) \longrightarrow J(B)$  are morphisms in  $\text{Dial}_2(\text{Sets})_!$ . Then we define the morphism  $(j_1, j_2) = ((f, g), \lambda j. \lambda w. F(h_1(j), w) \circ F(h_2(j), w)) : J(C) \longrightarrow J(A \times B)$ , where  $(f, g) = \lambda w. (f(w), g(w))$  is the unique morphism from the cartesian structure of  $\text{Sets}$ , and  $h_1(j) = \lambda u. \iota_1(j(u, g(w)))$  and  $h_2(j) = \lambda u. \iota_2(j(u, f(w)))$ . Notice that  $F$  is defined in terms of unique morphisms, and thus,  $(j_1, j_2)$  is unique.

All of the previous morphisms satisfy the usual diagram for cartesian products.

By Lemma 2 we know  $\text{Dial}_2(\text{Sets})$  is a full LNL model, and thus, there is an adjunction between  $\text{Dial}_2(\text{Sets})$  and  $\text{Dial}_2(\text{Sets})_!$  where the left-adjoint is the forgetful functor  $G : \text{Dial}_2(\text{Sets})_! \longrightarrow \text{Dial}_2(\text{Sets})$ , and the right adjoint is the free functor  $J : \text{Dial}_2(\text{Sets}) \longrightarrow \text{Dial}_2(\text{Sets})_!$ . Therefore,  $\text{Dial}_2(\text{Sets})$  is a model of full tensorial logic, as expected.

## 5.1 Double Negation Translation

In this section we show that we can use intuitionistic negation – which we showed was tensorial in the previous section – to define a negative translation of multi-conclusion linear logic (Figure 3) into FILL where implication plays a central role. Mellies and Tabareau give a negative translation of the multiplicative fragment of linear logic into tensorial logic [18] using tensor as the main connective. For example, they define  $(A \otimes B)^N = \neg(\neg(A)^N \otimes \neg(B)^N)$ , and thus, they simulate par using tensor and negation. This definition would cause some syntactic issues with FILL, because the left-rule to par requires the let-pattern term defined in Definition 3, thus, encoding par in terms of tensor would require the let-pattern term to be used in the left-rule for tensor. While simulating par, using tensor and negation, can be seen as useful, in applications where only the tensor product can be actually calculated, in other applications we do have an extra bifunctor like par. This is true in the case of FILL, so we can modify Mellies and Tabareau’s translation into one that better fits the source logical system.

The following definition defines a translation of linear logic formulas into FILL formulas.

**Definition 12.** *The following is the double-negation translation of linear logic into FILL:*



$$\begin{array}{c}
\frac{}{A \vdash A} \text{ LL-AX} \qquad \frac{\Gamma \vdash A, \Delta \quad \Gamma', A \vdash \Delta'}{\Gamma, \Gamma' \vdash \Delta, \Delta'} \text{ LL-CUT} \qquad \frac{\Gamma \vdash \Delta}{\Gamma, \top \vdash \Delta} \text{ LL-TL} \qquad \frac{}{\cdot \vdash \top} \text{ LL-TR} \\
\frac{\Gamma, A, B \vdash \Delta}{\Gamma, A \otimes B \vdash \Delta} \text{ LL-TENL} \qquad \frac{\Gamma \vdash A, \Delta \quad \Gamma' \vdash B, \Delta'}{\Gamma, \Gamma' \vdash A \otimes B, \Delta, \Delta'} \text{ LL-TENR} \qquad \frac{}{\perp \vdash \cdot} \text{ LL-PL} \qquad \frac{\Gamma \vdash \Delta}{\Gamma \vdash \perp, \Delta} \text{ LL-PR} \\
\frac{\Gamma, A \vdash \Delta \quad \Gamma', B \vdash \Delta'}{\Gamma, \Gamma', A \wp B \vdash \Delta, \Delta'} \text{ LL-PARL} \qquad \frac{\Gamma \vdash \Delta, A, B, \Delta'}{\Gamma \vdash \Delta, A \wp B, \Delta'} \text{ LL-PARR} \qquad \frac{\Gamma \vdash A, \Delta \quad \Gamma', B \vdash \Delta'}{\Gamma, A \multimap B, \Gamma' \vdash \Delta, \Delta'} \text{ LL-IMPL} \\
\frac{\Gamma, A \vdash B, \Delta}{\Gamma \vdash A \multimap B, \Delta} \text{ LL-IMPR} \qquad \frac{\Gamma, A, B \vdash \Delta}{\Gamma, B, A \vdash \Delta} \text{ LL-EXL} \qquad \frac{\Gamma \vdash \Delta_1, A, B, \Delta_2}{\Gamma \vdash \Delta_1, B, A, \Delta_2} \text{ LL-EXR}
\end{array}$$

**Fig. 3.** Multi-Conclusion Linear Logic

$$\begin{aligned}
(\top)^N &= \top \\
(\perp)^N &= \perp \\
(A^\perp)^N &= \neg((A)^N) \\
(A \wp B)^N &= \neg\neg((A)^N) \wp \neg\neg((B)^N) \\
(A \otimes B)^N &= \neg\neg((A)^N) \otimes \neg\neg((B)^N)
\end{aligned}$$

The main point of the previous definition is that because FILL has intuitionistic versions of all of the operators of linear logic we can give a very natural translation that preserves these operators by double negating their arguments.

At this point we need to extend the translation of linear logic formulas to sequents. However, we must be careful, because in FILL implication has the FILL restriction, and thus, if we choose the wrong translation then we will run into problems trying to satisfy the FILL condition. The method we employ here is to first translate a linear logic sequent into a single-sided sequent, and then translate that to FILL using the well-known translation. That is, it is easy to see that any linear logic sequent  $A_1, \dots, A_i \vdash B_1, \dots, B_j$  is logically equivalent to the sequent  $\cdot \vdash A_1^\perp, \dots, A_i^\perp, B_1, \dots, B_j$ . Then we translate the latter into FILL as  $x_1 : \neg((A_1^\perp)^N), \dots, x_i : \neg((A_i^\perp)^N), y_1 : \neg((B_1)^N), \dots, y_j : \neg((B_j)^N) \vdash \cdot$  for any free variables  $x_1, \dots, x_i$  and  $y_1, \dots, y_j$ , but this is equivalent to  $x_1 : \neg\neg((A_1)^N), \dots, x_i : \neg\neg((A_i)^N), y_1 : \neg((B_1)^N), \dots, y_j : \neg((B_j)^N) \vdash \cdot$ . The reader may realize that this is indeed the translation of single-sided classical linear logic into single-conclusion intuitionistic linear logic. This translation also has the benefit that we do not have to worry about mentioning terms in the statement of the result.

**Lemma 7 (Negative Translation).** *If  $A_1, \dots, A_i \vdash B_1, \dots, B_j$  is derivable, then for any unique fresh variables  $x_1, \dots, x_i$ , and  $y_1, \dots, y_j$ , the sequent  $x_1 : \neg\neg((A_1)^N), \dots, x_i : \neg\neg((A_i)^N), y_1 : \neg((B_1)^N), \dots, y_j : \neg((B_j)^N) \vdash \cdot$  is derivable.*

*Proof.* This is a proof by induction on the form of the assumed sequent  $A_1, \dots, A_i \vdash B_1, \dots, B_j$ . Throughout the proof if  $\Gamma = A_1, \dots, A_i$ , then  $\neg(\Gamma)^N = x_1 : \neg((A_1)^N), \dots, x_i : \neg((A_i)^N)$  for some unique fresh variables  $x_1, \dots, x_i$ . We will use this notation when applying the inductive hypothesis to make the presentation easier to read.

Case.

$$\frac{}{A \vdash A} \text{ Ax}$$

It suffices to derive the sequent  $x : \neg\neg((A)^N), y : \neg((A)^N) \vdash \cdot$  which is equivalent to the sequent  $x : \neg((A)^N) \multimap \perp, y : \neg((A)^N) \vdash \cdot$ . The latter is derivable as follows:

$$\frac{\frac{y : \neg((A)^N) \vdash y : \neg((A)^N)}{\text{Ax}} \quad \frac{}{w : \perp \vdash \cdot} \text{PL}}{x : \neg((A)^N) \multimap \perp, y : \neg((A)^N) \vdash \cdot} \text{IMPL}$$

Case.

$$\frac{\Gamma_1 \vdash A, \Delta_1 \quad \Gamma_2, A \vdash \Delta_2}{\Gamma_1, \Gamma_2 \vdash \Delta_1, \Delta_2} \text{CUT}$$

By the induction hypothesis we know the following:

$$\begin{aligned} & \neg(\neg((\Gamma_1)^N)), x : \neg((A)^N), \neg((\Delta_1)^N) \vdash \cdot \\ & \neg(\neg((\Gamma_2)^N)), y : \neg(\neg((A)^N)), \neg((\Delta_2)^N) \vdash \cdot \end{aligned}$$

It suffices to derive the sequent  $\neg(\neg((\Gamma_1)^N)), \neg(\neg((\Gamma_2)^N)), \neg((\Delta_1)^N), \neg((\Delta_2)^N) \vdash \cdot$  which we do as follows:

$$\frac{\frac{\frac{\neg(\neg((\Gamma_1)^N)), x : \neg((A)^N), \neg((\Delta_1)^N) \vdash \cdot}{\neg(\neg((\Gamma_1)^N)), x : \neg((A)^N), \neg((\Delta_1)^N) \vdash \circ : \perp} \text{PR}}{\neg(\neg((\Gamma_1)^N)), \neg((\Delta_1)^N) \vdash \lambda x. \circ : \neg(\neg((A)^N))} \text{IMPR} \quad \neg(\neg((\Gamma_2)^N)), y : \neg(\neg((A)^N)), \neg((\Delta_2)^N) \vdash \cdot}{\neg(\neg((\Gamma_1)^N)), \neg(\neg((\Gamma_2)^N)), \neg((\Delta_1)^N), \neg((\Delta_2)^N) \vdash \cdot} \text{CUT}$$

Case.

$$\frac{\Gamma \vdash \Delta}{\Gamma, \top \vdash \Delta} \text{TL}$$

First, note that  $(\top)^N = \top$ . By the induction hypothesis we know  $\neg(\neg((\Gamma)^N)), \neg((\Delta)^N) \vdash \cdot$ . We obtain our result by the following derivation:

$$\frac{\frac{\frac{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N) \vdash \cdot}{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), z : \top \vdash \cdot} \text{TL}}{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), z : \top \vdash t : \perp} \text{PR}}{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N) \vdash t : \neg \top} \text{IMPR} \quad \frac{}{y : \perp \vdash \cdot} \text{PL}}{\neg(\neg((\Gamma)^N)), x : \neg(\neg \top), \neg((\Delta)^N) \vdash \cdot} \text{IMPL}$$

Note that let  $x$  be  $*$  in  $\cdot = \cdot$  for any variable  $x$ .

Case.

$$\frac{}{\cdot \vdash \top} \text{TR}$$

It suffices to derive the sequent  $x : \neg((\top)^N) \vdash \cdot$ , but this is equivalent to the sequent  $x : \neg \top \vdash \cdot$  which is derivable as follows:

$$\frac{\frac{}{\cdot \vdash * : \top} \text{TR} \quad \frac{}{y : \perp \vdash \cdot} \text{PL}}{x : \neg \top \vdash \cdot} \text{IMPL}$$

Case.

$$\frac{\Gamma, A, B \vdash \Delta}{\Gamma, A \otimes B \vdash \Delta} \text{TENL}$$

By the induction hypothesis we have the following sequent:

$$\neg(\neg((\Gamma)^N)), x : \neg\neg((A)^N), y : \neg\neg((B)^N), \neg((\Delta)^N) \vdash \cdot$$

Our result follows from the following derivation:

$$\frac{\frac{\frac{\neg(\neg((\Gamma)^N)), w : \neg\neg((A)^N), v : \neg\neg((B)^N), \neg((\Delta)^N) \vdash \cdot}{\neg(\neg((\Gamma)^N)), w : \neg\neg((A)^N), v : \neg\neg((B)^N), \neg((\Delta)^N) \vdash \circ : \perp} \text{PR}}{\neg(\neg((\Gamma)^N)), x : \neg\neg((A)^N) \otimes \neg\neg((B)^N), \neg((\Delta)^N) \vdash \text{let } x \text{ be } (w \otimes v) \text{ in } \circ : \perp} \text{TENL}}{\frac{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N) \vdash \lambda x. \text{let } x \text{ be } (w \otimes v) \text{ in } \circ : \neg(\neg\neg((A)^N) \otimes \neg\neg((B)^N))}{\neg(\neg((\Gamma)^N)), z : \neg\neg(\neg\neg((A)^N) \otimes \neg\neg((B)^N)), \neg((\Delta)^N) \vdash \cdot} \text{IMPR} \quad \frac{}{y : \perp \vdash \cdot} \text{PL}} \text{IMPL}$$

Case.

$$\frac{\Gamma \vdash A, \Delta \quad \Gamma' \vdash B, \Delta'}{\Gamma, \Gamma' \vdash A \otimes B, \Delta, \Delta'} \text{TENR}$$

By the induction hypothesis we know the following:

$$\begin{aligned} &\neg(\neg((\Gamma)^N)), x : \neg\neg((A)^N), \neg((\Delta)^N) \vdash \cdot \\ &\neg(\neg((\Gamma')^N)), y : \neg\neg((B)^N), \neg((\Delta')^N) \vdash \cdot \end{aligned}$$

We obtain our result by the following derivation:

$$\frac{\frac{\frac{\neg(\neg((\Gamma)^N)), x : \neg\neg((A)^N), \neg((\Delta)^N) \vdash \cdot}{\neg(\neg((\Gamma)^N)), x : \neg\neg((A)^N), \neg((\Delta)^N) \vdash \circ : \perp} \text{PR}}{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N) \vdash \lambda x. \circ : \neg\neg((A)^N)} \text{IMPR} \quad \frac{\frac{\frac{\neg(\neg((\Gamma')^N)), y : \neg\neg((B)^N), \neg((\Delta')^N) \vdash \cdot}{\neg(\neg((\Gamma')^N)), y : \neg\neg((B)^N), \neg((\Delta')^N) \vdash \circ : \perp} \text{PR}}{\neg(\neg((\Gamma')^N)), \neg((\Delta')^N) \vdash \lambda y. \circ : \neg\neg((B)^N)} \text{IMPR}}{\frac{\neg(\neg((\Gamma)^N)), \neg(\neg((\Gamma')^N)), \neg((\Delta)^N), \neg((\Delta')^N) \vdash (\lambda x. \circ) \otimes (\lambda y. \circ) : \neg\neg((A)^N) \otimes \neg\neg((B)^N)}{\neg(\neg((\Gamma)^N)), \neg(\neg((\Gamma')^N)), z : \neg(\neg\neg((A)^N) \otimes \neg\neg((B)^N)), \neg((\Delta)^N), \neg((\Delta')^N) \vdash \cdot} \text{TENR} \quad \frac{}{w : \perp \vdash \cdot} \text{PL}} \text{IMPL}$$

Case.

$$\frac{}{\perp \vdash \cdot} \text{PL}$$

It suffices to show that  $z : \neg(\perp)^N \vdash \cdot$  is derivable, which is equivalent to showing that the sequent  $z : \neg\neg(\perp) \vdash \cdot$  is derivable. We obtain our result by the following derivation:

$$\frac{\frac{\frac{}{x : \perp \vdash \cdot} \text{PL}}{x : \perp \vdash \circ : \perp} \text{PR}}{\cdot \vdash \lambda x. \circ : \neg(\perp)} \text{IMPR} \quad \frac{}{w : \perp \vdash \cdot} \text{PL}}{z : \neg\neg(\perp) \vdash \cdot} \text{IMPL}$$

Case.

$$\frac{\Gamma \vdash \Delta}{\Gamma \vdash \perp, \Delta} \text{PR}$$

By the induction hypothesis we know the following sequent is derivable:

$$\neg(\neg((\Gamma)^N)), \neg((\Delta)^N) \vdash \cdot$$

It suffices to show that  $\neg(\neg((\Gamma)^N)), z : \neg(\perp)^N, \neg((\Delta)^N) \vdash \cdot$  which is equivalent to the sequent  $\neg(\neg((\Gamma)^N)), z : \neg(\perp), \neg((\Delta)^N) \vdash \cdot$ . The latter is derivable as follows:

$$\frac{\frac{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N) \vdash \cdot}{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N) \vdash \circ : \perp} \text{PR} \quad \frac{}{w : \perp \vdash \cdot} \text{PL}}{\neg(\neg((\Gamma)^N)), z : \neg(\perp), \neg((\Delta)^N) \vdash \cdot} \text{IMPL}$$

Case.

$$\frac{\Gamma, A \vdash \Delta \quad \Gamma', B \vdash \Delta'}{\Gamma, \Gamma', A \wp B \vdash \Delta, \Delta'} \text{PARL}$$

By the induction hypothesis we know the following:

$$\begin{aligned} &\neg(\neg((\Gamma)^N)), x : \neg\neg((A)^N), \neg((\Delta)^N) \vdash \cdot \\ &\neg(\neg((\Gamma')^N)), v : \neg\neg((B)^N), \neg((\Delta')^N) \vdash \cdot \end{aligned}$$

It suffices to show that the sequent

$$\neg(\neg((\Gamma)^N)), \neg(\neg((\Gamma')^N)), z : \neg\neg((A \wp B)^N), \neg((\Delta)^N), \neg((\Delta')^N) \vdash \cdot$$

is derivable which is equivalent to the sequent

$$\neg(\neg((\Gamma)^N)), \neg(\neg((\Gamma')^N)), z : \neg\neg(\neg\neg((A)^N) \wp \neg\neg((B)^N)), \neg((\Delta)^N), \neg((\Delta')^N) \vdash \cdot$$

The latter is derivable as follows:

$$\frac{D \quad \frac{}{w : \perp \vdash \cdot} \text{PL}}{\neg(\neg((\Gamma)^N)), \neg(\neg((\Gamma')^N)), z : \neg\neg(\neg\neg((A)^N) \wp \neg\neg((B)^N)), \neg((\Delta)^N), \neg((\Delta')^N) \vdash \cdot} \text{IMPL}$$

where

$D :$

$$\frac{\frac{\frac{\neg(\neg((\Gamma)^N)), x : \neg\neg((A)^N), \neg((\Delta)^N) \vdash \cdot \quad \neg(\neg((\Gamma')^N)), v : \neg\neg((B)^N), \neg((\Delta')^N) \vdash \cdot}{\neg(\neg((\Gamma)^N)), \neg(\neg((\Gamma')^N)), y : \neg\neg((A)^N) \wp \neg\neg((B)^N), \neg((\Delta)^N), \neg((\Delta')^N) \vdash \cdot} \text{PARL}}{\neg(\neg((\Gamma)^N)), \neg(\neg((\Gamma')^N)), y : \neg\neg((A)^N) \wp \neg\neg((B)^N), \neg((\Delta)^N), \neg((\Delta')^N) \vdash \circ : \perp} \text{PR}}{\neg(\neg((\Gamma)^N)), \neg(\neg((\Gamma')^N)), \neg((\Delta)^N), \neg((\Delta')^N) \vdash \lambda y. \circ : \neg(\neg\neg((A)^N) \wp \neg\neg((B)^N))} \text{IMPR}$$

Case.

$$\frac{\Gamma \vdash \Delta, A, B, \Delta'}{\Gamma \vdash \Delta, A \wp B, \Delta'} \text{PARR}$$

By the induction hypothesis we know the following is derivable:

$$\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), x : \neg((A)^N), y : \neg((B)^N), \neg((\Delta')^N) \vdash \cdot$$

It suffices to show that the sequent  $\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), z : \neg((A \wp B)^N), \neg((\Delta')^N)$  is derivable, but this sequent is equivalent to the sequent  $\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), z : \neg(\neg\neg((A)^N) \wp \neg\neg((B)^N)), \neg((\Delta')^N)$ . We derive the latter as follows:

$$\begin{array}{c}
\frac{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), x : \neg((A)^N), y : \neg((B)^N), \neg((\Delta')^N) \vdash \cdot}{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), x : \neg((A)^N), y : \neg((B)^N), \neg((\Delta')^N) \vdash \circ : \perp} \text{PR} \\
\frac{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), x : \neg((A)^N), \neg((\Delta')^N) \vdash (\lambda y. \circ) : \neg(\neg((B)^N))}{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), x : \neg((A)^N), \neg((\Delta')^N) \vdash \circ : \perp, (\lambda y. \circ) : \neg(\neg((B)^N))} \text{IMPR} \\
\frac{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), x : \neg((A)^N), \neg((\Delta')^N) \vdash (\lambda x. \circ) : \neg(\neg((A)^N)), (\lambda y. \circ) : \neg(\neg((B)^N))}{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), \neg((\Delta')^N) \vdash (\lambda x. \circ) \wp (\lambda y. \circ) : \neg(\neg((A)^N)) \wp \neg(\neg((B)^N))} \text{PARR} \\
\frac{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N), \neg((\Delta')^N) \vdash (\lambda x. \circ) \wp (\lambda y. \circ) : \neg(\neg((A)^N)) \wp \neg(\neg((B)^N)), \neg((\Delta')^N)}{w : \perp \vdash \cdot} \text{IMPL} \\
\text{PL}
\end{array}$$

Case.

$$\frac{\Gamma, A \vdash \Delta}{\Gamma \vdash A^\perp, \Delta} \text{NEGL}$$

By the induction hypothesis we know the following:

$$\neg(\neg((\Gamma)^N)), x : \neg(\neg((A)^N)), \neg((\Delta)^N) \vdash \cdot$$

It suffices to show that the sequent  $\neg(\neg((\Gamma)^N)), x : \neg((A^\perp)^N), \neg((\Delta)^N) \vdash \cdot$ , but this is equivalent to the sequent  $\neg(\neg((\Gamma)^N)), x : \neg(\neg((A)^N)), \neg((\Delta)^N) \vdash \cdot$ , but this is equivalent to the induction hypothesis.

Case.

$$\frac{\Gamma \vdash A, \Delta}{\Gamma, A^\perp \vdash \Delta} \text{NEGL}$$

By the induction hypothesis we know the following:

$$\neg(\neg((\Gamma)^N)), x : \neg((A)^N), \neg((\Delta)^N) \vdash \cdot$$

It suffices to show that the sequent  $\neg(\neg((\Gamma)^N)), z : \neg(\neg((A^\perp)^N)), \neg((\Delta)^N) \vdash \cdot$ , but this is equivalent to the sequent  $\neg(\neg((\Gamma)^N)), z : \neg(\neg(\neg((A)^N))), \neg((\Delta)^N) \vdash \cdot$ . The latter is derived as follows:

$$\frac{\frac{\frac{\neg(\neg((\Gamma)^N)), x : \neg((A)^N), \neg((\Delta)^N) \vdash \cdot}{\neg(\neg((\Gamma)^N)), x : \neg((A)^N), \neg((\Delta)^N) \vdash \circ : \perp} \text{PR}}{\neg(\neg((\Gamma)^N)), \neg((\Delta)^N) \vdash \lambda x. \circ : \neg(\neg((A)^N))} \text{IMPR}}{w : \perp \vdash \cdot} \text{IMPL} \\
\neg(\neg((\Gamma)^N)), z : \neg(\neg(\neg((A)^N))), \neg((\Delta)^N) \vdash \cdot$$

Case.

$$\frac{\Gamma, A, B \vdash \Delta}{\Gamma, B, A \vdash \Delta} \text{EXL}$$

This case follows by first applying the induction hypothesis and then applying the rule `EXL`.

Case.

$$\frac{\Gamma \vdash \Delta_1, A, B, \Delta_2}{\Gamma \vdash \Delta_1, B, A, \Delta_2} \text{EXR}$$

This case follows by first applying the induction hypothesis and then applying the rule `EXL`.

## 6 Conclusion and Future Work

We first recalled the definition of full intuitionistic linear logic using the left rule for par proposed by Bellin in Section 2, but using only proof-theoretic methods, no proof nets. We then directly proved cut-elimination for FILL in Section 3 by adapting the well-known cut-elimination procedure for classical linear logic to FILL.

In Section 4 we showed that the category  $\text{Dial}_2(\text{Sets})$ , a model of FILL, is a full LNL model by showing that it is a full linear category, and then replaying the proof that linear categories are LNL models by Benton. Then in Section 5 we showed that  $\text{Dial}_2(\text{Sets})$  is a model of full tensorial logic. The point of this exercise in categorical logic is to show that, despite linear logicians infatuation with linear negation, there is value in keeping all your connectives independent of each other. Only making them definable in terms of others, for specific applications.

Games, especially programming language games are the main motivation for Tensorial Logic and have been one of the sources of intuitions in linear logic all along. Since we are interested in the applications of tensorial logic to concurrency, we would like to see if our slightly more general framework can be applied to this task, just as well as tensorial logic.

Independently of the envisaged applications to programming, we are also interested in developing a “man in the street” game-like explanation for the finer-grained connectives of FILL, especially for par, the multiplicative disjunction. The second author has talked about games for FILL in the style of Lorenzen [10], building up on the work of Rahman [14,20]. Rahman showed that Lorenzen games could be defined for classical linear logic [19] and was able to define a sound and complete semantics in Lorenzen games for classical linear logic. While Rahman does mention that one could adopt a particular structural rule that enforces intuitionism, we have not seen a complete proof of soundness and completeness for this semantics. We plan to show that by adopting this rule we actually obtain a sound and complete semantics in Lorenzen games for FILL.

## References

1. Gianluigi Bellin. Subnets of proof-nets in multiplicative linear logic with mix. *Mathematical Structures in Computer Science*, 7:663–669, 12 1997.
2. P. N. Benton. A mixed linear and non-linear logic: Proofs, terms and models (preliminary report). Technical Report UCAM-CL-TR-352, University of Cambridge Computer Laboratory, 1994.
3. G. M. Bierman. *On Intuitionistic Linear Logic*. PhD thesis, Wolfson College, Cambridge, December 1993.
4. Gavin Bierman. A note on full intuitionistic linear logic. *Annals of Pure and Applied Logic*, 79(3):281 – 287, 1996.
5. Torben Braüner and Valeria Paiva. A formulation of linear logic based on dependency-relations. In Mogens Nielsen and Wolfgang Thomas, editors, *Computer Science Logic*, volume 1414 of *Lecture Notes in Computer Science*, pages 129–148. Springer Berlin Heidelberg, 1998.
6. R. Clouston, J. Dawson, R. Gore, and A. Tiu. Annotation-Free Sequent Calculi for Full Intuitionistic Linear Logic – Extended Version. *ArXiv e-prints*, July 2013.
7. J.R.B. Cockett and R.A.G. Seely. Weakly distributive categories. *Journal of Pure and Applied Algebra*, 114(2):133 – 173, 1997.
8. Valeria de Paiva. *The Dialectica Categories*. PhD thesis, University of Cambridge, 1988.
9. Valeria de Paiva. Dialectica categories. In J. Gray and A. Scedrov, editors, *Categories in Computer Science and Logic*, volume 92, pages 47–62. American Mathematical Society, 1989.
10. Valeria de Paiva. Abstract: Lorenzen games for full intuitionistic linear logic. *14th International Congress of Logic, Methodology and Philosophy of Science (CLMPS)*, 2011.
11. Valeria de Paiva and Luiz Carlos Pereira. A short note on intuitionistic propositional logic with multiple conclusions. *MANUSCRITO - Rev. Int. Fil.*, 28(2):317 – 329, jul-dez 2005.
12. A. G. Dragalin. *Mathematical Intuitionism. Introduction to Proof Theory*, volume 67 of *Translations of Mathematical Monographs*. American Mathematical Society, 1988.
13. Martin Hyland and Valeria de Paiva. Full intuitionistic linear logic (extended abstract). *Annals of Pure and Applied Logic*, 64(3):273 – 291, 1993.
14. Laurent Keiff. Dialogical logic. In Edward N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*. Summer 2011 edition, 2011.

15. S. Maehara. Eine darstellung der intuitionistischen logik in der klassischen. *Nagoya Mathematical Journal*, 7:45–64, 1954.
16. Paul-Andre Mellies. *Categorical Semantics of Linear Logic*. 2009.
17. Paul-André Melliés and Nicolas Tabareau. Linear continuations and duality. Technical report, Université Paris VII, 2008.
18. Paul-André Melliès and Nicolas Tabareau. Resource modalities in tensor logic. *Annals of Pure and Applied Logic*, 161(5):632 – 653, 2010.
19. S. Rahman. Un desafio para las teorías cognitivas de la competencia lógica: los fundamentos pragmáticos de la semántica de la lógica lineal. *Manuscrito*, XXV(2):381–432, October 2002.
20. Shahid Rahman and Laurent Keiff. On how to be a dialogician. In Daniel Vanderveken, editor, *Logic, Thought and Action*, volume 2 of *Logic, Epistemology, and the Unity of Science*, pages 359–408. Springer Netherlands, 2005.
21. Harold Schellinx. Some syntactical observations on linear logic. *Journal of Logic and Computation*, 1(4):537–559, 1991.
22. R. Seely. Linear logic, \*-autonomous categories and cofree coalgebras. In *Computer Science Logic*, 1989.
23. G. Takeuti. *Proof Theory*. Amsterdam: North-Holland, 1975.